

FORM PT01390  
(REV. 11-2000)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371

ATTORNEY'S DOCKET NUMBER

030639.0031.US1

U.S. APPLICATION NO. If known, sec 37 CFR 1.5

09/889331

INTERNATIONAL APPLICATION NO.

PCT/US00/00942

INTERNATIONAL FILING DATE

14 January 2000 (14.01.00)

PRIORITY DATE CLAIMED

14 January 1999 (14.01.99)

TITLE OF INVENTION METHODS FOR GLUCAGON SUPPRESSION

APPLICANT(S) FOR DO/EO/US YOUNG, Andrew; GEDULIN, Bronislava

Applicant herewith submits to the United States Designate/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below.
4. ☒ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
  - a. ☐ is attached hereto (required only if not communicated by the International Bureau).
  - b. ☐ has been communicated by the International Bureau.
  - c. ☒ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
  - a. ☐ is attached hereto.
  - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
7. ☐ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
  - a. ☐ are attached hereto (required only if not communicated by the International Bureau).
  - b. ☐ have been communicated by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☐ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)).
9. ☐ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

## Items 11 to 20 below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☐ A **FIRST** preliminary amendment.
14. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
15. ☐ A substitute specification.
16. ☐ A change of power of attorney and/or address letter.
17. ☐ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
18. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
19. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
20. ☐ Other items or information:



METHODS FOR GLUCAGON SUPPRESSIONRELATED APPLICATIONS

This application claims priority from U.S. Provisional Application 60/116,380, entitled "Novel Exendin Agonist Formulations And Methods Of Administration Thereof," filed January 14, 1999 (and the corresponding PCT application filed January 14, 2000, Serial No. [not yet assigned]), U.S. Provisional Application 60/132,017, entitled "Methods for Glucagon Suppression," filed April 30, 1999, and U.S. Provisional Application 60/[not yet assigned], entitled "Use of Exendins and Agonists Thereof for Modulation of Triglyceride Levels and Treatment of Dyslipidemia," filed January 10, 2000, the contents of which are hereby incorporated by reference in their entireties.

FIELD OF THE INVENTION

The present invention relates to methods of suppressing and/or lowering glucagon in a subject, comprising the administration of an exendin, an exendin agonist, or a modified exendin or exendin agonist having an exendin or exendin agonist peptide linked to one or more polyethylene glycol polymers or other compound useful to decrease renal clearance of the parent peptide. Such methods are useful, for example, in the treatment of hyperglucagonemia and other conditions in which lower levels of glucagon or suppression of glucagon secretion are of benefit.

BACKGROUND

The following description includes information that may be useful in understanding the present invention. It is not an admission that any of the information provided herein is prior art to the presently claimed invention, or relevant, nor that any of the publications specifically or implicitly referenced are prior art.

The extendins are peptides that are found in the salivary secretions of the Gila monster and the Mexican Beaded Lizard, reptiles that are endogenous to Arizona and Northern Mexico. Exendin-3 [SEQ. ID. NO. 1: His Ser Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro Ser-NH<sub>2</sub>] is present in the salivary secretions of *Heloderma horridum* (Mexican Beaded Lizard), and exendin-4 [SEQ. ID. NO. 2: His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro Ser-NH<sub>2</sub>] is present in the salivary secretions of *Heloderma suspectum* (Gila monster) (Eng, J., et al., J. Biol. Chem., 265:20259-62, 1990; Eng, J., et al., J. Biol. Chem., 267:7402-05, 1992). The amino acid sequence of exendin-3 is shown in Figure 1. The amino acid sequence of exendin-4 is shown in Figure 2. Exendin-4 was first thought to be a (potentially toxic) component of the venom. It now appears that exendin-4 is devoid of toxicity, and that it instead is made in salivary glands in the Gila monster.

The extendins have some sequence similarity to several members of the glucagon-like peptide family, with the highest homology, 53%, being to GLP-1[7-36]NH<sub>2</sub> [SEQ. ID. NO. 3] (Goke, et al., J. Biol. Chem., 268:19650-55, 1993). GLP-1[7-36]NH<sub>2</sub>, also sometimes referred to as proglucagon[78-107] or simply "GLP-1" as used most often herein, has an insulinotropic effect, stimulating insulin secretion from pancreatic beta-cells; GLP-1 has also been reported to inhibit glucagon secretion from pancreatic alpha-cells (Ørsov, et al., Diabetes, 42:658-61, 1993; D'Alessio, et al., J. Clin. Invest., 97:133-38, 1996). GLP-1 has been reported to inhibit gastric emptying (Willms B, et al., J Clin Endocrinol Metab 81 (1): 327-32, 1996; Wettergren A, et

al., Dig Dis Sci 38 (4): 665-73, 1993), and gastric acid secretion (Schjoldager BT, et al., Dig Dis Sci 34 (5): 703-8, 1989; O'Halloran DJ, et al., J Endocrinol 126 (1): 169-73, 1990; Wettergren A, et al., Dig Dis Sci 38 (4): 665-73, 1993)). GLP-1[7-37], which has an additional glycine residue at its carboxy terminus, is reported to stimulate insulin secretion in humans (Ørskov, et al., Diabetes, 42:658-61, 1993). A transmembrane G-protein adenylate-cyclase-coupled receptor said to be responsible at least in part for the insulinotropic effect of GLP-1 has reportedly been cloned from a beta-cell line (Thorens, Proc. Natl. Acad. Sci. USA 89:8641-45, 1992). GLP-1 has been the focus of significant investigation in recent years due to its reported action on the amplification of stimulated insulin production (Byrne MM, Goke B. Lessons from human studies with glucagon-like peptide-1: Potential of the gut hormone for clinical use. In: Fehmann HC, Goke B. Insulinotropic Gut Hormone Glucagon-Like Peptide 1. Basel, Switzerland: Karger, 1997:219-33).

Other reports relate to the inhibition of gastric emptying (Wettergren A, et al., Truncated GLP-1 (proglucagon 78-107-amide) inhibits gastric and pancreatic functions in man, Dig. Dis. Sci. 1993 Apr;38(4):665-73), inhibition of glucagon secretion (Creutzfeldt WOC, et al., Glucagonostatic actions and reduction of fasting hyperglycemia by exogenous glucagon-like peptide I(7-36) amide in type I diabetic patients, Diabetes Care 1996;19(6):580-6), and a purported role in appetite control (Turton MD, et al., A role for glucagon-like peptide-1 in the central regulation of feeding, Nature 1996 Jan;379(6560):69-72).

GLP-1 has also been reported to restore islet glucose sensitivity in aging rats, restoring their glucose tolerance to that of younger rats (Egan JM, et al., Glucagon-like

249/167

peptide-1 restores acute-phase insulin release to aged rats, *Diabetologia* 1997 Jun;40(Suppl 1):A130). However, the short duration of biological action of GLP-1 *in vivo* is one feature of the peptide that has hampered its development as a therapeutic agent. Various methods have been tried to prolong the half-life of GLP-1 or GLP-1(7-37), including attempts to alter their amino acid sequence and to deliver them using certain formulations (see, e.g., European Patent Application, entitled "Prolonged Delivery of Peptides," by Darley, et al., publication number 0 619 322 A2, regarding the inclusion of polyethylene glycol in formulations containing GLP-1 (7-37)).

Pharmacological studies have led to reports that exendin-4 can act at GLP-1 receptors on certain insulin-secreting cells, at dispersed acinar cells from guinea pig pancreas, and at parietal cells from stomach; the peptide is also reported to stimulate somatostatin release and inhibit gastrin release in isolated stomachs (Goke, et al., J. Biol. Chem. 268:19650-55, 1993; Schepp, et al., Eur. J. Pharmacol., 69:183-91, 1994; Eissele, et al., Life Sci., 55:629-34, 1994). Exendin-3 and exendin-4 were reportedly found to stimulate cAMP production in, and amylase release from, pancreatic acinar cells (Malhotra, R., et al., Regulatory Peptides, 41:149-56, 1992; Raufman, et al., J. Biol. Chem. 267:21432-37, 1992; Singh, et al., Regul. Pept. 53:47-59, 1994). Additionally, exendin-4 has a significantly longer duration of action than GLP-1. For example, in one experiment, glucose lowering by exendin-4 in diabetic mice was reported to persist for several hours, and, depending on dose, for up to 24 hours (Eng J. Prolonged effect of exendin-4 on hyperglycemia of db/db mice, Diabetes 1996 May; 45(Suppl 2):152A (abstract 554)). Based on their insulinotropic activities, the use of exendin-3 and exendin-

4 for the treatment of diabetes mellitus and the prevention of hyperglycemia has been proposed (Eng, U.S. Patent No. 5,424,286).

5 The results of an investigation of whether exendins are the species homolog of mammalian GLP-1 was reported by Chen and Drucker who cloned the exendin gene from the Gila monster (J. Biol. Chem. 272(7):4108-15 (1997)). The observation that the Gila monster also has separate genes for proglucagons (from which GLP-1 is processed), that are  
10 more similar to mammalian proglucagon than exendin, indicates that exendins are not merely species homologs of GLP-1.

To date, agents that serve to delay gastric emptying have generally found a place in medicine as diagnostic aids  
15 in gastrointestinal radiological examinations. For example, glucagon is a polypeptide hormone that is produced by the alpha cells of the pancreatic islets of Langerhans. It is a hyperglycemic agent that mobilizes glucose by activating hepatic glycogenolysis. It can to a lesser extent stimulate  
20 the secretion of pancreatic insulin. Glucagon is used in the treatment of insulin-induced hypoglycemia, for example, when administration of glucose intravenously is not possible. However, as glucagon reduces the motility of the gastro-intestinal tract it is also used as a diagnostic aid  
25 in gastrointestinal radiological examinations. Glucagon has also been used in several studies to treat various painful gastrointestinal disorders associated with spasm. Daniel, et al. (Br. Med. J., 3:720, 1974) reported quicker symptomatic relief of acute diverticulitis in patients  
30 treated with glucagon compared with those who had been treated with analgesics or antispasmodics. A review by Glauser, et al. (J. Am. Coll. Emergency Physns, 8:228, 1979) described relief of acute esophageal food obstruction

following glucagon therapy. In another study, glucagon significantly relieved pain and tenderness in 21 patients with biliary tract disease compared with 22 patients treated with placebo (M.J. Stower, et al., Br. J. Surg., 69:591-2, 1982).

Methods for regulating gastrointestinal motility using amylin agonists are described in commonly owned International Application No. PCT/US94/10225, published March 16, 1995.

Methods for regulating gastrointestinal motility using exendin agonists are described in commonly owned U.S. Patent Application Serial No. 08/908,867, filed August 8, 1997 entitled "Methods for Regulating Gastrointestinal Motility," which application is a continuation-in-part of U.S. Patent Application Serial No. 08/694,954, filed August 8, 1996.

Methods for reducing food intake using exendin agonists are described in commonly owned U.S. Patent Application Serial No. 09/003,869, filed January 7, 1998, entitled "Use of Exendin and Agonists Thereof for the Reduction of Food Intake," which claims the benefit of U.S. Provisional Application Nos. 60/034,905 filed January 7, 1997, 60/055,404 filed August 7, 1997, 60/065,442 filed November 14, 1997 and 60/066,029 filed November 14, 1997.

Novel exendin agonist compounds are described in commonly owned PCT Application Serial No. PCT/US98/16387 filed August 6, 1998, entitled "Novel Exendin Agonist Compounds," which claims the benefit of U.S. Patent Application Serial No. 60/055,404, filed August 8, 1997.

Other novel exendin agonists are described in commonly owned PCT Application Serial No. PCT/US98/24210, filed November 13, 1998, entitled "Novel Exendin Agonist Compounds," which claims the benefit of U.S. Provisional Application No. 60/065,442 filed November 14, 1997.



Still other novel exendin agonists are described in commonly owned PCT Application Serial No. PCT/US98/24273, filed November 13, 1998, entitled "Novel Exendin Agonist Compounds," which claims the benefit of U.S. Provisional Application No. 60/066,029 filed November 14, 1997.

Other recent advances in exendin related technology are described in U.S. Provisional Patent Application Serial No. 60/075,122, filed February 13, 1998, entitled "Inotropic and Diuretic Effects of Exendin and GLP-1" and in U.S.

Provisional Patent Application Serial No. 60/116,380, filed January 14, 1998, entitled "Novel Exendin Agonist Formulations and Methods of Administration Thereof".

Polyethylene glycol (PEG) modification of therapeutic peptides and proteins may yield both advantages and disadvantages. While PEG modification may lead to improved circulation time, reduced antigenicity and immunogenicity, improved solubility, resistance to proteolysis, improved bioavailability, reduced toxicity, improved stability, and easier formulation of peptides (See, Francis et al., International Journal of Hematology, 68:1-18, 1998) problems with PEGylation in most cases is substantial reduction in bioactivity. Id. In addition, most methods involve use of linkers that have several types of adverse effects including immunogenicity, instability, toxicity, and reactivity. Id.

Glucagonoma (tumor of glucagon-secreting cells) produces, in addition to glucose intolerance, a skin condition, necrolytic migratory erythema. This is a raised scaly red rash, sometimes blistering and eventually crusting, localized to the face, abdomen, extremities and perineum. It can also be associated with inflammation of the tongue and mouth, and diseased nails and thinning of the hair. The condition is reported to respond to octreotide, a glucagonostatic hormone analog. The compounds described

herein are also useful as glucagonastatic agents and thus in the treatment of this disease, which was first described in 1966 (Kaplan, L.M. Endocrine Tumors of the Gastrointestinal Tract and Pancreas. Ch 262, p1392: In Harrison's Principles of Internal Medicine, 12th Edition. McGraw-Hill Inc, New York, 1991). The compounds described herein that are useful for lowering glucagon levels and/or suppressing glucagon secretion include exendin, exendin agonists, and modified exendins and exendin agonists and related formulations, and dosage formulations.

The contents of the above-identified articles, patents, and patent applications, and all other documents mentioned or cited herein, are hereby incorporated by reference in their entirety. Applicants reserve the right to physically incorporate into this application any and all materials and information from any such articles, patents, patent applications, or other documents mentioned or cited herein.

#### SUMMARY OF THE INVENTION

The present invention relates to methods for lowering glucagon levels and/or suppressing glucagon secretion in a subject. It also relates to the treatment of hyperglucagonemia and conditions that benefit from administration of glucagonostatic agents, including but not limited to necrolytic migratory erythema.

Thus, in one aspect, the invention relates to the use of an exendin, an exendin agonist, or a modified exendin or exendin agonist having an exendin or exendin agonist linked to one or more polyethylene glycol polymers, or other molecular weight enhancing molecules, for lowering glucagon levels in a subject.

In another aspect, the invention relates to the use of an exendin, an exendin agonist, or a modified exendin or

exendin agonist having an exendin or exendin agonist linked to one or more polyethylene glycol polymers or other compounds useful to decrease renal clearance of the parent peptide, for suppressing glucagon secretion in a subject.

5 In still another aspect, the invention relates to the use of an exendin, an exendin agonist, or a modified exendin or exendin agonist having an exendin or exendin agonist linked to one or more polyethylene glycol polymers, or other molecular weight enhancing molecules, for treating  
10 conditions associated with hyperglucagonemia.

In yet another aspect, the invention relates to the use of an exendin, an exendin agonist, or a modified exendin or exendin agonist having an exendin or exendin agonist linked to one or more polyethylene glycol polymers, or other  
15 molecular weight enhancing molecules, for treating a subject with a glucagonoma or necrolytic migratory erythema.

In preferred embodiments, the exendin is exendin-4. In other preferred embodiments, the modified exendin or exendin agonist has a molecular weight that is greater than the  
20 molecular weight of the exendin or exendin agonist (preferably about 10%, 50% or 90% greater), the modified exendin or exendin agonist has a negative charge that is greater than the negative charge of the exendin or exendin agonist (preferably about 10%, 50% or 90% greater), the  
25 modified exendin or exendin agonist has a kidney clearance that is less than the kidney clearance of the exendin or exendin agonist (preferably about 10%, 50% or 90% less), the modified exendin or exendin agonist has a half-life that is greater than the half-life of the exendin or exendin agonist  
30 (preferably about 10%, 50% or 90% greater), the modified exendin or exendin agonist has a immunogenicity/antigenicity that is less than the immunogenicity/antigenicity of the exendin or exendin agonist, the modified exendin or exendin

agonist has a solubility that is greater than the solubility of the exendin or exendin agonist (preferably about 10%, 50% or 90% greater), the modified exendin or exendin agonist has a proteolysis rate that is less than the proteolysis rate of the exendin or exendin agonist (preferably about 10%, 50% or 90% less), the modified exendin or exendin agonist has a toxicity that is less than the toxicity of the exendin or exendin agonist, the modified exendin or exendin agonist has a stability that is greater than the stability of the exendin or exendin agonist, and the modified exendin or exendin agonist has a permeability/biological function that is greater or less than the permeability/biological function of the exendin or exendin agonist (preferably about 10%, 50% or 90% greater or less).

The exendin or exendin agonist may be linked to one, two or three polyethylene glycol polymers. The polyethylene glycol polymers may preferably have molecular weights between 500 and 20,000. In a preferred embodiment, the modified exendin or exendin agonist is one of compounds 201-217, more preferably one of compounds 209, 210 and 213, or one of compounds 201 and 202, or one of compounds 216 and 217 (See Example 4 below).

The polyethylene glycol polymers are preferably linked to an amino, carboxyl, or thio group, and may be linked by N or C termini of side chains of lysine, aspartic acid, glutamic acid, or cysteine, or alternatively, the polyethylene glycol polymers may be linked with diamine and dicarboxylic groups. The exendin or exendin agonist is preferably linked to the polyethylene glycol polymers through an epsilon amino group on a lysine amino acid of the exendin or exendin agonist.

By "exendin agonist" is meant a compound which mimics the effects of exendins, e.g., on gastric motility and

gastric emptying (namely, a compound which effectively binds to the receptor at which exendins exert their action on gastric motility and gastric emptying, preferably an analog or derivative of an exendin) or a compound, e.g., that mimics the effects of exendin on the reduction of food intake by binding to the receptor or receptors where exendin causes this effect. Preferred exendin agonist compounds include those described in United States Patent Application Serial No. 90/003,869, entitled, "Use of Exendin And Agonists Thereof For The Reduction of Food Intake", filed January 7, 1998, (and the priority applications thereto) which enjoys common ownership with the present application and which is incorporated by this reference into the present application as though fully set forth herein. Effects of exendins or exendin agonists can be identified, evaluated, or screened for, using the methods described herein, or other methods known in the art for determining exendin effects.

In another aspect, a therapeutically effective amount of an amylin agonist is also administered to the subject. In a preferred aspect, the amylin agonist is an amylin or an amylin agonist analog such as <sup>25,28,29</sup>Pro-human-amylin. (also known as "pramlintide," and previously referred to as "AC-137" and described in "Amylin Agonist Peptides and Uses Therefor," U.S. Patent No. 5,686,511, issued November 11, 1997), or salmon calcitonin.

Preferably, the subject is a vertebrate, more preferably a mammal, and most preferably a human. In preferred aspects, the exendin, exendin agonist, or modified exendin or exendin agonist of the invention is administered parenterally, more preferably by injection. In a most preferred aspect, the injection is a peripheral injection. Preferably, about 1 µg-30 µg to about 5 mg of the modified exendin or exendin agonist of the invention is administered per day. More

preferably, about 1-30  $\mu$ g to about 2mg, or about 1-30  $\mu$ g to about 1mg of the modified exendin or exendin agonist of the invention is administered per day. Most preferably, about 3  $\mu$ g to about 500  $\mu$ g of the modified exendin or exendin agonist of the invention is administered per day.

Preferred exendins or exendin agonists for modification and use include:

exendin-4 (1-30) [SEQ ID NO 4: His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly];

exendin-4 (1-30) amide [SEQ ID NO 5: His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly-NH<sub>2</sub>];

exendin-4 (1-28) amide [SEQ ID NO 6: His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn-NH<sub>2</sub>];

<sup>14</sup>Leu, <sup>25</sup>Phe exendin-4 amide [SEQ ID NO 7: His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn Gly Gly Pro Ser Ser Gly Ala Pro Pro Pro Ser-NH<sub>2</sub>];

<sup>14</sup>Leu, <sup>25</sup>Phe exendin-4 (1-28) amide [SEQ ID NO 8: His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>]; and

<sup>14</sup>Leu, <sup>22</sup>Ala, <sup>25</sup>Phe exendin-4 (1-28) amide [SEQ ID NO 9: His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu Ala Ile Glu Phe Leu Lys Asn-NH<sub>2</sub>].

### Definitions

In accordance with the present invention and as used herein, the following terms are defined to have the following meanings, unless explicitly stated otherwise.


The term "amino acid" refers to natural amino acids, unnatural amino acids, and amino acid analogs, all in their

D and L stereoisomers if their structure allow such stereoisomeric forms. Natural amino acids include alanine (Ala), arginine (Arg), asparagine (Asn), aspartic acid (Asp), cysteine (Cys), glutamine (Gln), glutamic acid (Glu), glycine (Gly), histidine (His), isoleucine (Ile), leucine (Leu), Lysine (Lys), methionine (Met), phenylalanine (Phe), proline (Pro), serine (Ser), threonine (Thr), typtophan (Trp), tyrosine (Tyr) and valine (Val). Unnatural amino acids include, but are not limited to azetidinecarboxylic acid, 2-aminoadipic acid, 3-aminoadipic acid, beta-alanine, aminopropionic acid, 2-aminobutyric acid, 4-aminobutyric acid, 6-aminocaproic acid, 2-aminoheptanoic acid, 2-aminoisobutyric acid, 3-aminoisbutyric acid, 2-aminopimelic acid, tertiary-butylglycine, 2,4-diaminoisobutyric acid, desmosine, 2,2'-diaminopimelic acid, 2,3-diaminopropionic acid, N-ethylglycine, N-ethylasparagine, homoproline, hydroxylysine, allo-hydroxylysine, 3-hydroxyproline, 4-hydroxyproline, isodesmosine, allo-isoleucine, N-methylalanine, N-methylglycine, N-methylisoleucine, N-methylpentylglycine, N-methylvaline, naphthalanine, norvaline, norleucine, ornithine, pentylglycine, pipecolic acid and thioproline. Amino acid analogs include the natural and unnatural amino acids which are chemically blocked, reversibly or irreversibly, or modified on their N-terminal amino group or their side-chain groups, as for example, methionine sulfoxide, methionine sulfone, S-(carboxymethyl)-cysteine, S-(carboxymethyl)-cysteine sulfoxide and S-(carboxymethyl)-cysteine sulfone.

The term "amino acid analog" refers to an amino acid wherein either the C-terminal carboxy group, the N-terminal amino group or side-chain functional group has been chemically codified to another functional group. For example, aspartic acid-(beta-methyl ester) is an amino acid

analog of aspartic acid; N-ethylglycine is an amino acid analog of glycine; or alanine carboxamide is an amino acid analog of alanine.

The term "amino acid residue" refers to radicals having the structure: (1)  $-C(O)-R-NH-$ , wherein R typically is  $-CH(R')-$ , wherein R' is an amino acid side chain, typically H or a carbon containing substituent;



or (2) , wherein p is 1, 2 or 3 representing the azetidinecarboxylic acid, proline or pipecolic acid residues, respectively.

The term "lower" referred to herein in connection with organic radicals such as alkyl groups defines such groups with up to and including about 6, preferably up to and including 4 and advantageously one or two carbon atoms. Such groups may be straight chain or branched chain.

"Pharmaceutically acceptable salt" includes salts of the compounds of the present invention derived from the combination of such compounds and an organic or inorganic acid. In practice the use of the salt form amounts to use of the base form. The compounds of the present invention are useful in both free base and salt form, with both forms being considered as being within the scope of the present invention.

In addition, the following abbreviations stand for the following:

"ACN" or "CH<sub>3</sub>CN" refers to acetonitrile.

"Boc", "tBoc" or "Tboc" refers to t-butoxy carbonyl.

"DCC" refers to N,N'-dicyclohexylcarbodiimide.

"Fmoc" refers to fluorenylmethoxycarbonyl.

"HBTU" refers to 2-(1H-benzotriazol-1-yl)-



1,1,3,3,-tetramethyluronium hexafluorophosphate.

"HOBt" refers to 1-hydroxybenzotriazole monohydrate.

"homop" or hPro" refers to homoproline.

"MeAla" or "Nme" refers to N-methylalanine.

5 "naph" refers to naphthylalanine.

"pG" or pGly" refers to pentylglycine.

"tBuG" refers to tertiary-butylglycine.

"ThioP" or tPro" refers to thioproline.

"3Hyp" refers to 3-hydroxyproline

10 "4Hyp" refers to 4-hydroxyproline

"NAG" refers to N-alkylglycine

"NAPG" refers to N-alkylpentylglycine

"Norval" refers to norvaline

"Norleu" refers to norleucine

15 Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

#### BRIEF DESCRIPTION OF THE DRAWINGS

20 Figure 1 depicts the amino acid sequence for exendin-3 [SEQ. ID. NO. 1].

Figure 2 depicts the amino acid sequence for exendin-4 [SEQ. ID. NO. 2].

25 Figure 3 depicts the amino acid sequences for certain exendin agonist compounds useful in the present invention [SEQ. ID. NOS. 10 TO 40].

Figure 4 depicts the amino acid sequences for certain compounds of the present invention, Compounds 1-174.

30 Figure 5 is a graph showing the effect of functional nephrectomy on exendin-4 clearance.

Figure 6 is a graph showing the terminal decay of exendin-4 plasma levels in nephrectomized and sham subjects.

DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to relates to methods of suppressing and/or lowering glucagon in a subject, comprising the administration of an exendin, an exendin  
5 agonist, or a modified exendin or exendin agonist having an exendin or exendin agonist peptide linked to one or more polyethylene glycol polymers or other compound useful to increase molecular weight. Such methods are useful, for example, in the treatment of hyperglucagonemia and other  
10 conditions in which lower levels of glucagon or suppression of glucagon secretion are of benefit. Such conditions include, but are not limited to, glucagonoma and necrolytic migratory erythema.

15 Modified Exendins And Exendin Agonists

The modified exendins and exendin agonists of the present invention include, for example, one or more PEG polymers linked to an exendin or exendin agonist, such as a  
20 naturally occurring exendin, a synthetic exendin or an exendin agonist.

Exendin-4

Exendin-4 is a naturally occurring peptide isolated from the salivary secretions of the Gila monster. Animal  
25 testing of exendin-4 has shown that its ability to lower blood glucose persists for several hours. Exendin-4, a 39-amino acid polypeptide, is synthesized using solid phase synthesis as described herein, and this synthetic material has been shown to be identical to that of native exendin-4.

30 As described herein, the nonclinical pharmacology of exendin-4 has been studied. In the brain, exendin-4 binds principally to the area *postrema* and *nucleus tractus solitarius* region in the hindbrain and to the subfornical

organ in the forebrain. Exendin-4 binding has been observed in the rat and mouse brain and kidney. The structures to which exendin-4 binds in the kidney are unknown.

Various experiments have compared the biologic actions of exendin-4 and GLP-1 and demonstrated a more favorable spectrum of properties for exendin-4. A single subcutaneous dose of exendin-4 lowered plasma glucose in *db/db* (diabetic) and *ob/ob* (diabetic obese) mice by up to 40%. In Diabetic Fatty Zucker (ZDF) rats, 5 weeks of treatment with exendin-4 lowered HbA<sub>1c</sub> (a measure of glycosylated hemoglobin used to evaluate plasma glucose levels) by up to 41%. Insulin sensitivity was also improved by 76% following 5 weeks of treatment in obese ZDF rats. In glucose intolerant primates, dose-dependent decreases in plasma glucose were also observed.

An insulinitropic action of exendin-4 has also been observed in rodents, improving insulin response to glucose by over 100% in non-fasted Harlan Sprague Dawley (HSD) rats, and by up to -10-fold in non-fasted *db/db* mice. Higher pretreatment plasma glucose concentrations were associated with greater glucose-lowering effects. Thus the observed glucose lowering effect of exendin-4 appears to be glucose-dependent, and minimal if animals are already euglycemic.

Exendin-4 dose dependently slowed gastric emptying in HSD rats and was -90-fold more potent than GLP-1 for this action. Exendin-4 has also been shown to reduce food intake in NIH/Sw (Swiss) mice following peripheral administration, and was at least 1000 times more potent than GLP-1 for this action. Exendin-4 reduced plasma glucagon concentrations by approximately 40% in anesthetized ZDF rats during hyperinsulinemic, hyperglycemic clamp conditions, but did not affect plasma glucagon concentrations during euglycemic conditions in normal rats. Exendin-4 has been shown to

dose-dependently reduce body weight in obese ZDF rats, while in lean ZDF rats, the observed decrease in body weight appears to be transient.

Through effects on lowering glucagon and suppressing glucagon secretion, exendins, exendin agonists, and modified exendins or exendin agonists containing exendin-4, for example, will be useful in people who would benefit from lowered glucagon, for example, people with glucagonoma and necrolytic migratory erythema, and people with diabetes whether or not they retain the ability to secrete insulin. See Example 5.

The toxicology of exendin-4 has been investigated in single-dose studies in mice, rats and monkeys, repeated-dose (up to 28 consecutive daily doses) studies in rats and monkeys and *in vitro* tests for mutagenicity and chromosomal alterations. To date, no deaths have occurred, and there have been no observed treatment-related changes in hematology, clinical chemistry, or gross or microscopic tissue changes. Exendin-4 was demonstrated to be non-mutagenic, and did not cause chromosomal aberrations at the concentrations tested (up to 5000 µg/mL).

In support of the investigation of the nonclinical pharmacokinetics and metabolism of exendin-4, a number of immunoassays have been developed. A radioimmunoassay with limited sensitivity (~100 pM) was used in initial pharmacokinetic studies. A two-site IRMA assay for exendin-4 was subsequently validated with a lower limit of quantitation of 15 pM. The bioavailability of exendin-4, given subcutaneously, was found to be approximately 50-80% using the radioimmunoassay. This was similar to that seen following intraperitoneal administration (48-60%). Peak plasma concentrations ( $C_{max}$ ) occurred between 30 and 43 minutes ( $T_{max}$ ). Both  $C_{max}$  and AUC values were monotonically

related to dose. The apparent terminal half-life for exendin-4 given subcutaneously was approximately 90-110 minutes. This was significantly longer than the 14-41 minutes seen following intravenous dosing. Similar results were obtained using the IRMA assay. Degradation studies with exendin-4 compared to GLP-1 indicate that exendin-4 is relatively resistant to degradation.

#### Exendin Agonists

The structure activity relationship (SAR) of exendin was investigated for structures that may relate to the antidiabetic activity of exendin, for its stability to metabolism, and for improvement of its physical characteristics, especially as it pertains to peptide stability and to amenability to alternative delivery systems, and various exendin agonist peptide compounds have been invented. Exendin agonists include exendin peptide analogs in which one or more naturally occurring amino acids are eliminated or replaced with another amino acid(s). Preferred exendin agonists are agonist analogs of exendin-4. Particularly preferred exendin agonists include those described in commonly owned PCT Application Serial No. PCT/US98/16387 filed August 6, 1998, entitled "Novel Exendin Agonist Compounds," which claims the benefit of U.S. Patent Application Serial No. 60/055,404, filed August 8, 1997; commonly owned PCT Application Serial No. PCT/US98/24210, filed November 13, 1998, entitled "Novel Exendin Agonist Compounds," which claims the benefit of U.S. Provisional Application No. 60/065,442 filed November 14, 1997; and, commonly owned PCT Application Serial No. PCT/US98/24273, filed November 13, 1998, entitled "Novel Exendin Agonist Compounds," which claims the benefit of U.S. Provisional Application No. 60/066,029 filed November 14, 1997, all of

which are incorporated herein by reference in their entirety, including any drawings.

Activity as exendin agonists can be indicated, for example, by activity in the assays described below. Effects of exendins or exendin agonists on gastric motility and gastric emptying can be identified, evaluated, or screened for, using the methods described herein, or other art-known or equivalent methods for determining gastric motility. Negative receptor assays or screens for exendin agonist compounds or candidate exendin agonist compounds, such as an amylin receptor assay/screen using an amylin receptor preparation as described in U.S. Patent No. 5,264,372, issued November 23, 1993, the contents of which are incorporated herein by reference, one or more calcitonin receptor assays/screens using, for example, T47D and MCF7 breast carcinoma cells, which contain calcium receptors coupled to the stimulation of adenylyl cyclase activity, and/or a CGRP receptor assay/screen using, for example, SK-N-MC cells.

One such method for use in identifying or evaluating the ability of a compound to slow gastric motility, involves: (a) bringing together a test sample and a test system, the test sample containing one or more test compounds, the test system containing a system for evaluating gastric motility, the system being characterized in that it exhibits, for example, elevated plasma glucose in response to the introduction to the system of glucose or a meal; and, (b) determining the presence or amount of a rise in plasma glucose in the system. Positive and/or negative controls may be used as well.

Also included within the scope of the present invention are pharmaceutically acceptable salts of the compounds of

formula (I-VIII) and pharmaceutical compositions including said compounds and salts thereof.

FORMULA I

- 5        Exendin agonist compounds also include those described in U.S. Provisional Application No. 60/065,442, including compounds of the formula (I) [SEQ ID NO. 41]:

Xaa<sub>1</sub> Xaa<sub>2</sub> Xaa<sub>3</sub> Gly Xaa<sub>5</sub> Xaa<sub>6</sub> Xaa<sub>7</sub> Xaa<sub>8</sub> Xaa<sub>9</sub> Xaa<sub>10</sub>

Xaa<sub>11</sub> Xaa<sub>12</sub> Xaa<sub>13</sub> Xaa<sub>14</sub> Xaa<sub>15</sub> Xaa<sub>16</sub> Xaa<sub>17</sub> Ala Xaa<sub>19</sub> Xaa<sub>20</sub>

- 10    Xaa<sub>21</sub> Xaa<sub>22</sub> Xaa<sub>23</sub> Xaa<sub>24</sub> Xaa<sub>25</sub> Xaa<sub>26</sub> Xaa<sub>27</sub> Xaa<sub>28</sub>-Z<sub>1</sub>; wherein

Xaa<sub>1</sub> is His, Arg or Tyr;

Xaa<sub>2</sub> is Ser, Gly, Ala or Thr;

Xaa<sub>3</sub> is Asp or Glu;

- 15    Xaa<sub>5</sub> is Ala or Thr;

Xaa<sub>6</sub> is Ala, Phe, Tyr or naphthylalanine;

Xaa<sub>7</sub> is Thr or Ser;

Xaa<sub>8</sub> is Ala, Ser or Thr;

Xaa<sub>9</sub> is Asp or Glu;

- 20    Xaa<sub>10</sub> is Ala, Leu, Ile, Val, pentylglycine or Met;

Xaa<sub>11</sub> is Ala or Ser;

Xaa<sub>12</sub> is Ala or Lys;

Xaa<sub>13</sub> is Ala or Gln;

Xaa<sub>14</sub> is Ala, Leu, Ile, pentylglycine, Val or Met;

- 25    Xaa<sub>15</sub> is Ala or Glu;

Xaa<sub>16</sub> is Ala or Glu;

Xaa<sub>17</sub> is Ala or Glu;

Xaa<sub>19</sub> is Ala or Val;

Xaa<sub>20</sub> is Ala or Arg;

- 30    Xaa<sub>21</sub> is Ala or Leu;

Xaa<sub>22</sub> is Ala, Phe, Tyr or naphthylalanine;

Xaa<sub>23</sub> is Ile, Val, Leu, pentylglycine, tert-butylglycine or Met;

Xaa<sub>24</sub> is Ala, Glu or Asp;

Xaa<sub>25</sub> is Ala, Trp, Phe, Tyr or naphthylalanine;

Xaa<sub>26</sub> is Ala or Leu;

Xaa<sub>27</sub> is Ala or Lys;

5 Xaa<sub>28</sub> is Ala or Asn;

Z<sub>1</sub> is -OH,

-NH<sub>2</sub>

Gly-Z<sub>2</sub>,

Gly Gly-Z<sub>2</sub>,

10 Gly Gly Xaa<sub>31</sub>-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala-Z<sub>2</sub>,

15 Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub>-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub>-Z<sub>2</sub> or

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub>-Z<sub>2</sub>;

Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently Pro,  
homoproline, 3Hyp, 4Hyp, thioproline,

20 N-alkylglycine, N-alkylpentylglycine or

N-alkylalanine; and

Z<sub>2</sub> is -OH or -NH<sub>2</sub>;

provided that no more than three of Xaa<sub>3</sub>, Xaa<sub>5</sub>, Xaa<sub>6</sub>, Xaa<sub>8</sub>,

Xaa<sub>10</sub>, Xaa<sub>11</sub>, Xaa<sub>12</sub>, Xaa<sub>13</sub>, Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>, Xaa<sub>17</sub>, Xaa<sub>19</sub>,

25 Xaa<sub>20</sub>, Xaa<sub>21</sub>, Xaa<sub>24</sub>, Xaa<sub>25</sub>, Xaa<sub>26</sub>, Xaa<sub>27</sub> and Xaa<sub>28</sub> are Ala.

Preferred N-alkyl groups for N-alkylglycine, N-  
alkylpentylglycine and N-alkylalanine include lower alkyl  
groups preferably of 1 to about 6 carbon atoms, more  
preferably of 1 to 4 carbon atoms.

30 Preferred exendin agonist compounds include those  
wherein Xaa<sub>1</sub> is His or Tyr. More preferably Xaa<sub>1</sub> is His.

Preferred are those compounds wherein Xaa<sub>2</sub> is Gly.



Preferred are those compounds wherein Xaa<sub>14</sub> is Leu, pentylglycine or Met.

Preferred compounds are those wherein Xaa<sub>25</sub> is Trp or Phe.

5 Preferred compounds are those where Xaa<sub>6</sub> is Phe or naphthylalanine; Xaa<sub>22</sub> is Phe or naphthylalanine and Xaa<sub>23</sub> is Ile or Val.

Preferred are compounds wherein Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently selected from Pro, homoproline, 10 thioproline and N-alkylalanine.

Preferably Z<sub>1</sub> is -NH<sub>2</sub>.

Preferably Z<sub>2</sub> is -NH<sub>2</sub>.

According to one aspect, preferred are compounds of formula (I) wherein Xaa<sub>1</sub> is His or Tyr, more preferably His; 15 Xaa<sub>2</sub> is Gly; Xaa<sub>6</sub> is Phe or naphthylalanine; Xaa<sub>14</sub> is Leu, pentylglycine or Met; Xaa<sub>22</sub> is Phe or naphthylalanine; Xaa<sub>23</sub> is Ile or Val; Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently selected from Pro, homoproline, thioproline or N-alkylalanine. More preferably Z<sub>1</sub> is -NH<sub>2</sub>.

20 According to an especially preferred aspect, especially preferred compounds include those of formula (I) wherein: Xaa<sub>1</sub> is His or Arg; Xaa<sub>2</sub> is Gly or Ala; Xaa<sub>3</sub> is Asp or Glu; Xaa<sub>5</sub> is Ala or Thr; Xaa<sub>6</sub> is Ala, Phe or naphthylalanine; Xaa<sub>7</sub> is Thr or Ser; Xaa<sub>8</sub> is Ala, Ser or Thr; Xaa<sub>9</sub> is Asp or Glu; 25 Xaa<sub>10</sub> is Ala, Leu or pentylglycine; Xaa<sub>11</sub> is Ala or Ser; Xaa<sub>12</sub> is Ala or Lys; Xaa<sub>13</sub> is Ala or Gln; Xaa<sub>14</sub> is Ala, Leu or pentylglycine; Xaa<sub>15</sub> is Ala or Glu; Xaa<sub>16</sub> is Ala or Glu; Xaa<sub>17</sub> is Ala or Glu; Xaa<sub>19</sub> is Ala or Val; Xaa<sub>20</sub> is Ala or Arg; Xaa<sub>21</sub> is Ala or Leu; Xaa<sub>22</sub> is Phe or naphthylalanine; Xaa<sub>23</sub> is Ile, 30 Val or tert-butylglycine; Xaa<sub>24</sub> is Ala, Glu or Asp; Xaa<sub>25</sub> is Ala, Trp or Phe; Xaa<sub>26</sub> is Ala or Leu; Xaa<sub>27</sub> is Ala or Lys; Xaa<sub>28</sub> is Ala or Asn; Z<sub>1</sub> is -OH, -NH<sub>2</sub>, Gly-Z<sub>2</sub>, Gly Gly-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub>-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub>-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub>-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub>-Z<sub>2</sub>; Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> being independently  
5 Pro homoproline, thioproline or N-methylalanine; and Z<sub>2</sub> being -OH or -NH<sub>2</sub>; provided that no more than three of Xaa<sub>3</sub>, Xaa<sub>5</sub>, Xaa<sub>6</sub>, Xaa<sub>8</sub>, Xaa<sub>10</sub>, Xaa<sub>11</sub>, Xaa<sub>12</sub>, Xaa<sub>13</sub>, Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>, Xaa<sub>17</sub>, Xaa<sub>19</sub>, Xaa<sub>20</sub>, Xaa<sub>21</sub>, Xaa<sub>24</sub>, Xaa<sub>25</sub>, Xaa<sub>26</sub>, Xaa<sub>27</sub> and Xaa<sub>28</sub> are Ala. Especially preferred compounds include those set  
10 forth in PCT application Serial No. PCT/US98/24210, filed November 13, 1998, entitled "Novel Exendin Agonist Compounds" identified therein as compounds 2-23.

According to an especially preferred aspect, provided are compounds where Xaa<sub>14</sub> is Leu, Ile, Val or pentylglycine,  
15 more preferably Leu or pentylglycine, and Xaa<sub>25</sub> is Phe, Tyr or naphthylalanine, more preferably Phe or naphthylalanine. These compounds will be less susceptible to oxidative degradation, both in vitro and in vivo, as well as during synthesis of the compound.

## FORMULA II

Exendin agonist compounds also include those described in U.S. Provisional Application No. 60/066,029, including compounds of the formula (II) [SEQ ID NO. 42]:

25 Xaa<sub>1</sub> Xaa<sub>2</sub> Xaa<sub>3</sub> Xaa<sub>4</sub> Xaa<sub>5</sub> Xaa<sub>6</sub> Xaa<sub>7</sub> Xaa<sub>8</sub> Xaa<sub>9</sub> Xaa<sub>10</sub>  
Xaa<sub>11</sub> Xaa<sub>12</sub> Xaa<sub>13</sub> Xaa<sub>14</sub> Xaa<sub>15</sub> Xaa<sub>16</sub> Xaa<sub>17</sub> Ala Xaa<sub>19</sub> Xaa<sub>20</sub>  
Xaa<sub>21</sub> Xaa<sub>22</sub> Xaa<sub>23</sub> Xaa<sub>24</sub> Xaa<sub>25</sub> Xaa<sub>26</sub> Xaa<sub>27</sub> Xaa<sub>28</sub>-Z<sub>1</sub>; wherein

Xaa<sub>1</sub> is His, Arg, Tyr, Ala, Norval, Val  
30 or Norleu;

Xaa<sub>2</sub> is Ser, Gly, Ala or Thr;

Xaa<sub>3</sub> is Ala, Asp or Glu;

Xaa<sub>4</sub> is Ala, Norval, Val, Norleu or Gly;

- Xaa<sub>5</sub> is Ala or Thr;  
Xaa<sub>6</sub> is Phe, Tyr or naphthylalanine;  
Xaa<sub>7</sub> is Thr or Ser;  
Xaa<sub>8</sub> is Ala, Ser or Thr;  
5 Xaa<sub>9</sub> is Ala, Norval, Val, Norleu, Asp or Glu;  
Xaa<sub>10</sub> is Ala, Leu, Ile, Val, pentylglycine or Met;  
Xaa<sub>11</sub> is Ala or Ser;  
Xaa<sub>12</sub> is Ala or Lys;  
Xaa<sub>13</sub> is Ala or Gln;  
10 Xaa<sub>14</sub> is Ala, Leu, Ile, pentylglycine, Val or Met;  
Xaa<sub>15</sub> is Ala or Glu;  
Xaa<sub>16</sub> is Ala or Glu;  
Xaa<sub>17</sub> is Ala or Glu;  
Xaa<sub>19</sub> is Ala or Val;  
15 Xaa<sub>20</sub> is Ala or Arg;  
Xaa<sub>21</sub> is Ala or Leu;  
Xaa<sub>22</sub> is Phe, Tyr or naphthylalanine;  
Xaa<sub>23</sub> is Ile, Val, Leu, pentylglycine, tert-butylglycine or Met;  
20 Xaa<sub>24</sub> is Ala, Glu or Asp;  
Xaa<sub>25</sub> is Ala, Trp, Phe, Tyr or naphthylalanine;  
Xaa<sub>26</sub> is Ala or Leu;  
Xaa<sub>27</sub> is Ala or Lys;  
Xaa<sub>28</sub> is Ala or Asn;  
25 Z<sub>1</sub> is -OH,  
-NH<sub>2</sub>,  
Gly-Z<sub>2</sub>,  
Gly Gly-Z<sub>2</sub>,  
Gly Gly Xaa<sub>31</sub>-Z<sub>2</sub>,  
30 Gly Gly Xaa<sub>31</sub> Ser-Z<sub>2</sub>,  
Gly Gly Xaa<sub>31</sub> Ser Ser-Z<sub>2</sub>,  
Gly Gly Xaa<sub>31</sub> Ser Ser Gly-Z<sub>2</sub>,  
Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub>-Z<sub>2</sub>,  
Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub>-Z<sub>2</sub>,  
Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub>-Z<sub>2</sub> or  
Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub> Xaa<sub>39</sub>-Z<sub>2</sub>;

5 wherein

Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently  
Pro, homoproline, 3Hyp, 4Hyp, thioproline,  
N-alkylglycine, N-alkylpentylglycine or  
N-alkylalanine; and

10 Z<sub>2</sub> is -OH or -NH<sub>2</sub>;

provided that no more than three of Xaa<sub>3</sub>, Xaa<sub>4</sub>, Xaa<sub>5</sub>, Xaa<sub>6</sub>,  
Xaa<sub>8</sub>, Xaa<sub>9</sub>, Xaa<sub>10</sub>, Xaa<sub>11</sub>, Xaa<sub>12</sub>, Xaa<sub>13</sub>, Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>,  
Xaa<sub>17</sub>, Xaa<sub>19</sub>, Xaa<sub>20</sub>, Xaa<sub>21</sub>, Xaa<sub>24</sub>, Xaa<sub>25</sub>, Xaa<sub>26</sub>, Xaa<sub>27</sub> and Xaa<sub>28</sub>  
are Ala; and provided also that, if Xaa<sub>1</sub> is His, Arg or Tyr,  
15 then at least one of Xaa<sub>3</sub>, Xaa<sub>4</sub> and Xaa<sub>9</sub> is Ala.

Preferred N-alkyl groups for N-alkylglycine, N-  
alkylpentylglycine and N-alkylalanine include lower alkyl  
groups preferably of 1 to about 6 carbon atoms, more  
preferably of 1 to 4 carbon atoms. Suitable compounds of  
20 formula (II) include those described in application Serial  
No. PCT/US98/24273, filed November 13, 1998, entitled "Novel  
Exendin Agonist Compounds", identified therein in Examples  
1-89 ("Compounds 1-89," respectively), as well as those  
corresponding compounds identified therein in Examples 104  
25 and 105.

Preferred such exendin agonist compounds include those  
wherein Xaa<sub>1</sub> is His, Ala or Norval. More preferably Xaa<sub>1</sub> is  
His or Ala. Most preferably Xaa<sub>1</sub> is His.

Preferred are those compounds of formula (II) wherein  
30 Xaa<sub>2</sub> is Gly.

Preferred are those compounds of formula (II) wherein  
Xaa<sub>3</sub> is Ala.

Preferred are those compounds of formula (II) wherein Xaa<sub>4</sub> is Ala.

Preferred are those compounds of formula (II) wherein Xaa<sub>9</sub> is Ala.

5 Preferred are those compounds of formula (II) wherein Xaa<sub>14</sub> is Leu, pentylglycine or Met.

Preferred compounds of formula (II) are those wherein Xaa<sub>25</sub> is Trp or Phe.

10 Preferred compounds of formula (II) are those where Xaa<sub>6</sub> is Ala, Phe or naphthylalanine; Xaa<sub>22</sub> is Phe or naphthylalanine; and Xaa<sub>23</sub> is Ile or Val.

Preferred are compounds of formula (II) wherein Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently selected from Pro, homoproline, thioproline and N-alkylalanine.

15 Preferably Z<sub>1</sub> is -NH<sub>2</sub>.

Preferably Z<sub>2</sub> is -NH<sub>2</sub>.

According to one aspect, preferred are compounds of formula (II) wherein Xaa<sub>1</sub> is Ala, His or Tyr, more preferably Ala or His; Xaa<sub>2</sub> is Ala or Gly; Xaa<sub>6</sub> is Phe or naphthylalanine; Xaa<sub>14</sub> is Ala, Leu, pentylglycine or Met; Xaa<sub>22</sub> is Phe or naphthylalanine; Xaa<sub>23</sub> is Ile or Val; Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently selected from Pro, homoproline, thioproline or N-alkylalanine; and Xaa<sub>39</sub> is Ser or Tyr, more preferably Ser. More preferably Z<sub>1</sub> is -NH<sub>2</sub>.

25 According to an especially preferred aspect, especially preferred compounds include those of formula (II) wherein: Xaa<sub>1</sub> is His or Ala; Xaa<sub>2</sub> is Gly or Ala; Xaa<sub>3</sub> is Ala, Asp or Glu; Xaa<sub>4</sub> is Ala or Gly; Xaa<sub>5</sub> is Ala or Thr; Xaa<sub>6</sub> is Phe or naphthylalanine; Xaa<sub>7</sub> is Thr or Ser; Xaa<sub>8</sub> is Ala, Ser or Thr; Xaa<sub>9</sub> is Ala, Asp or Glu; Xaa<sub>10</sub> is Ala, Leu or pentylglycine; Xaa<sub>11</sub> is Ala or Ser; Xaa<sub>12</sub> is Ala or Lys; Xaa<sub>13</sub> is Ala or Gln; Xaa<sub>14</sub> is Ala, Leu, Met or pentylglycine; Xaa<sub>15</sub> is Ala or Glu; Xaa<sub>16</sub> is Ala or Glu; Xaa<sub>17</sub> is Ala or Glu; Xaa<sub>19</sub> is Ala or Val;

Xaa<sub>20</sub> is Ala or Arg; Xaa<sub>21</sub> is Ala or Leu; Xaa<sub>22</sub> is Phe or naphthylalanine; Xaa<sub>23</sub> is Ile, Val or tert-butylglycine; Xaa<sub>24</sub> is Ala, Glu or Asp; Xaa<sub>25</sub> is Ala, Trp or Phe; Xaa<sub>26</sub> is Ala or Leu; Xaa<sub>27</sub> is Ala or Lys; Xaa<sub>28</sub> is Ala or Asn; Z<sub>1</sub> is -OH, -NH<sub>2</sub>, Gly-Z<sub>2</sub>, Gly Gly-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub>-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser Gly-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub>-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub>-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub>-Z<sub>2</sub> or Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub> Xaa<sub>39</sub>-Z<sub>2</sub>; Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> being independently Pro homoproline, thioproline or N-methylalanine; and Z<sub>2</sub> being -OH or -NH<sub>2</sub>; provided that no more than three of Xaa<sub>3</sub>, Xaa<sub>5</sub>, Xaa<sub>6</sub>, Xaa<sub>8</sub>, Xaa<sub>10</sub>, Xaa<sub>11</sub>, Xaa<sub>12</sub>, Xaa<sub>13</sub>, Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>, Xaa<sub>17</sub>, Xaa<sub>19</sub>, Xaa<sub>20</sub>, Xaa<sub>21</sub>, Xaa<sub>24</sub>, Xaa<sub>25</sub>, Xaa<sub>26</sub>, Xaa<sub>27</sub> and Xaa<sub>28</sub> are Ala; and provided also that, if Xaa<sub>1</sub> is His, Arg or Tyr, then at least one of Xaa<sub>3</sub>, Xaa<sub>4</sub> and Xaa<sub>9</sub> is Ala. Especially preferred compounds of formula (II) include those described in application Serial No. PCT/US98/24273, filed November 13, 1998, entitled "Novel Exendin Agonist Compounds" as having the amino acid sequence of SEQ. ID. NOS. 5-93 therein.

According to an especially preferred aspect, provided are compounds of formula (II) where Xaa<sub>14</sub> is Ala, Leu, Ile, Val or pentylglycine, more preferably Leu or pentylglycine, and Xaa<sub>25</sub> is Ala, Phe, Tyr or naphthylalanine, more preferably Phe or naphthylalanine. These compounds will be less susceptible to oxidative degradation, both in vitro and in vivo, as well as during synthesis of the compound.

### 30 FORMULA III

Also within the scope of the present invention are narrower genera of compounds having peptides of various lengths, for example genera of compounds which do not

include peptides having a length of 28, 29 or 30 amino acid residues, respectively. Additionally, the present invention includes narrower genera of compounds described in PCT application Serial No. PCT/US98/24210, filed November 13, 1998, entitled "Novel Exendin Agonist Compounds" and having particular amino acid sequences, for example, compounds of the formula (III) [SEQ. ID. NO. 43]:

Xaa<sub>1</sub> Xaa<sub>2</sub> Xaa<sub>3</sub> Gly Xaa<sub>5</sub> Xaa<sub>6</sub> Xaa<sub>7</sub> Xaa<sub>8</sub> Xaa<sub>9</sub> Xaa<sub>10</sub>  
10 Xaa<sub>11</sub> Xaa<sub>12</sub> Xaa<sub>13</sub> Xaa<sub>14</sub> Xaa<sub>15</sub> Xaa<sub>16</sub> Xaa<sub>17</sub> Ala Xaa<sub>18</sub> Xaa<sub>19</sub>  
Xaa<sub>20</sub> Xaa<sub>21</sub> Xaa<sub>22</sub> Xaa<sub>23</sub> Xaa<sub>24</sub> Xaa<sub>25</sub> Xaa<sub>26</sub> Xaa<sub>27</sub> Xaa<sub>28</sub>-Z<sub>1</sub>;

wherein

- Xaa<sub>1</sub> is His or Arg;  
15 Xaa<sub>2</sub> is Gly or Ala;  
Xaa<sub>3</sub> is Asp or Glu;  
Xaa<sub>5</sub> is Ala or Thr;  
Xaa<sub>6</sub> is Ala, Phe or naphthylalanine;  
Xaa<sub>7</sub> is Thr or Ser;  
20 Xaa<sub>8</sub> is Ala, Ser or Thr;  
Xaa<sub>9</sub> is Asp or Glu;  
Xaa<sub>10</sub> is Ala, Leu or pentylglycine;  
Xaa<sub>11</sub> is Ala or Ser;  
Xaa<sub>12</sub> is Ala or Lys;  
25 Xaa<sub>13</sub> is Ala or Gln;  
Xaa<sub>14</sub> is Ala, Leu or pentylglycine;  
Xaa<sub>15</sub> is Ala or Glu;  
Xaa<sub>16</sub> is Ala or Glu;  
Xaa<sub>17</sub> is Ala or Glu;  
30 Xaa<sub>19</sub> is Ala or Val;  
Xaa<sub>20</sub> is Ala or Arg;  
Xaa<sub>21</sub> is Ala or Leu;  
Xaa<sub>22</sub> is Phe or naphthylalanine;

Xaa<sub>23</sub> is Ile, Val or tert-butylglycine;

Xaa<sub>24</sub> is Ala, Glu or Asp;

Xaa<sub>25</sub> is Ala, Trp, or Phe;

Xaa<sub>26</sub> is Ala or Leu;

5 Xaa<sub>27</sub> is Ala or Lys;

Xaa<sub>28</sub> is Ala or Asn;

Z<sub>1</sub> is -OH,

-NH<sub>2</sub>,

Gly-Z<sub>2</sub>,

10 Gly Gly -Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub>-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly-Z<sub>2</sub>,

15 Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub>-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub>-Z<sub>2</sub> or Gly Gly  
Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub>-Z<sub>2</sub>;

20 Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently selected  
from the group consisting of Pro, homoproline,  
thioprolin and N-methylalanine; and

Z<sub>2</sub> is -OH or -NH<sub>2</sub>;

provided that no more than three of Xaa<sub>3</sub>, Xaa<sub>5</sub>, Xaa<sub>6</sub>, Xaa<sub>8</sub>,  
Xaa<sub>10</sub>, Xaa<sub>11</sub>, Xaa<sub>12</sub>, Xaa<sub>13</sub>, Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>, Xaa<sub>17</sub>, Xaa<sub>19</sub>,

25 Xaa<sub>20</sub>, Xaa<sub>21</sub>, Xaa<sub>24</sub>, Xaa<sub>25</sub>, Xaa<sub>26</sub>, Xaa<sub>27</sub> and Xaa<sub>28</sub> are Ala; and  
pharmaceutically acceptable salts thereof.

#### FORMULA IV

30 Additionally, the present invention includes narrower  
genera of peptide compounds described in PCT Application  
Serial No. PCT/US98/24273, filed November 13, 1998, entitled  
"Novel Exendin Agonist Compounds" as having particular amino  
acid sequences, for example, compounds of the formula [IV]



[SEQ. ID. NO. 44]:

Xaa<sub>1</sub> Xaa<sub>2</sub> Xaa<sub>3</sub> Xaa<sub>4</sub> Xaa<sub>5</sub> Xaa<sub>6</sub> Xaa<sub>7</sub> Xaa<sub>8</sub> Xaa<sub>9</sub> Xaa<sub>10</sub> Xaa<sub>11</sub> Xaa<sub>12</sub>  
Xaa<sub>13</sub> Xaa<sub>14</sub> Xaa<sub>15</sub> Xaa<sub>16</sub> Xaa<sub>17</sub> Ala Xaa<sub>18</sub> Xaa<sub>19</sub> Xaa<sub>20</sub> Xaa<sub>21</sub> Xaa<sub>22</sub> Xaa<sub>23</sub>  
5 Xaa<sub>24</sub> Xaa<sub>25</sub> Xaa<sub>26</sub> Xaa<sub>27</sub> Xaa<sub>28</sub>-Z<sub>1</sub>; wherein

- Xaa<sub>1</sub> is His or Ala;  
Xaa<sub>2</sub> is Gly or Ala;  
Xaa<sub>3</sub> is Ala, Asp or Glu;  
10 Xaa<sub>4</sub> is Ala or Gly;  
Xaa<sub>5</sub> is Ala or Thr;  
Xaa<sub>6</sub> is Phe or naphthylalanine;  
Xaa<sub>7</sub> is Thr or Ser;  
Xaa<sub>8</sub> is Ala, Ser or Thr;  
15 Xaa<sub>9</sub> is Ala, Asp or Glu;  
Xaa<sub>10</sub> is Ala, Leu or pentylglycine;  
Xaa<sub>11</sub> is Ala or Ser;  
Xaa<sub>12</sub> is Ala or Lys;  
Xaa<sub>13</sub> is Ala or Gln;  
20 Xaa<sub>14</sub> is Ala, Leu, Met or pentylglycine;  
Xaa<sub>15</sub> is Ala or Glu;  
Xaa<sub>16</sub> is Ala or Glu;  
Xaa<sub>17</sub> is Ala or Glu;  
Xaa<sub>19</sub> is Ala or Val;  
25 Xaa<sub>20</sub> is Ala or Arg;  
Xaa<sub>21</sub> is Ala or Leu;  
Xaa<sub>22</sub> is Phe or naphthylalanine;  
Xaa<sub>23</sub> is Ile, Val or tert-butylglycine;  
Xaa<sub>24</sub> is Ala, Glu or Asp;  
30 Xaa<sub>25</sub> is Ala, Trp or Phe;  
Xaa<sub>26</sub> is Ala or Leu;  
Xaa<sub>27</sub> is Ala or Lys;  
Xaa<sub>28</sub> is Ala or Asn;

Z<sub>1</sub> is -OH,

-NH<sub>2</sub>,

Gly-Z<sub>2</sub>,

Gly Gly-Z<sub>2</sub>

5 Gly Gly Xaa<sub>31</sub>-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala-Z<sub>2</sub>,

10 Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub>-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub>-Z<sub>2</sub>

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub>-Z<sub>2</sub>

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub>

Ser-Z<sub>2</sub>;

15 Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently Pro,  
homoproline, thioproline, or

N-methylalalanine; and

Z<sub>2</sub> is -OH or -NH<sub>2</sub>;

provided that no more than three of Xaa<sub>3</sub>, Xaa<sub>5</sub>, Xaa<sub>6</sub>, Xaa<sub>8</sub>,

20 Xaa<sub>10</sub>, Xaa<sub>11</sub>, Xaa<sub>12</sub>, Xaa<sub>13</sub>, Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>, Xaa<sub>17</sub>, Xaa<sub>19</sub>,

Xaa<sub>20</sub>, Xaa<sub>21</sub>, Xaa<sub>24</sub>, Xaa<sub>25</sub>, Xaa<sub>26</sub>, Xaa<sub>27</sub>, and Xaa<sub>28</sub> are Ala; and

provided that, if Xaa<sub>1</sub> is His, Arg or Tyr, then at least one  
of Xaa<sub>3</sub>, Xaa<sub>4</sub> and Xaa<sub>9</sub> is Ala; and pharmaceutically  
acceptable salts thereof.

25 Preferred compounds of formula (IV) include those  
wherein Xaa<sub>1</sub> is His, Ala, Norval or 4-imidazopropionyl.  
Preferably, Xaa<sub>1</sub> is His, or 4-imidazopropionyl or Ala, more  
preferably His or 4-imidazopropionyl.

30 Preferred compounds of formula (IV) include those  
wherein Xaa<sub>2</sub> is Gly.

Preferred compounds of formula (IV) include those  
wherein Xaa<sub>4</sub> is Ala.

Preferred compounds of formula (IV) include those

wherein Xaa<sub>9</sub> is Ala.

Preferred compounds of formula (IV) include those wherein Xaa<sub>14</sub> is Leu, pentylglycine or Met.

5 Preferred compounds of formula (IV) include those wherein Xaa<sub>25</sub> is Trp or Phe.

Preferred compounds of formula (IV) include those wherein Xaa<sub>6</sub> is Ala, Phe or naphthylalanine; Xaa<sub>22</sub> is Phe or naphthylalanine; and Xaa<sub>23</sub> is Ile or Val.

10 Preferred compounds of formula (IV) include those wherein Z<sub>1</sub> is -NH<sub>2</sub>.

Preferred compounds of formula (IV) include those wherein Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently selected from the group consisting of Pro, homoproline, thioproline and N-alkylalanine.

15 Preferred compounds of formula (IV) include those wherein Xaa<sub>39</sub> is Ser or Tyr, preferably Ser.

Preferred compounds of formula (IV) include those wherein Z<sub>2</sub> is -NH<sub>2</sub>.

20 Preferred compounds of formula (IV) include those wherein Z<sub>1</sub> is -NH<sub>2</sub>.

Preferred compounds of formula (IV) include those wherein Xaa<sub>21</sub> is Lys-NH<sup>c</sup>-R where R is Lys, Arg, C<sub>1</sub>-C<sub>10</sub> straight chain or branched alkanoyl.

25 Preferred compounds of formula (IV) include those wherein X<sub>1</sub> is Lys Asn, Lys-NH<sup>c</sup>-R Asn, or Lys-NH<sup>c</sup>-R Ala where R is Lys, Arg, C<sub>1</sub>-C<sub>10</sub> straight chain or branched alkanoyl.

Preferred compounds of formula (IV) include those having an amino acid sequence described in PCT application Serial No. PCT/US98/24273, filed November 13, 1998, entitled "Novel  
30 Exendin Agonist Compounds" as being selected from SEQ. ID. NOS. 95-110 therein.

FORMULA V

Also provided are compounds described in PCT application PCT/US98/24210, filed November 13, 1998, entitled "Novel Exendin Agonist Compounds", including  
5 compounds of the formula (V) [SEQ. ID. NO. 45]:

5

10

Xaa<sub>1</sub> Xaa<sub>2</sub> Xaa<sub>3</sub> Gly Xaa<sub>5</sub> Xaa<sub>6</sub> Xaa<sub>7</sub> Xaa<sub>8</sub> Xaa<sub>9</sub> Xaa<sub>10</sub>  
Xaa<sub>11</sub> Xaa<sub>12</sub> Xaa<sub>13</sub> Xaa<sub>14</sub> Xaa<sub>15</sub> Xaa<sub>16</sub> Xaa<sub>17</sub> Ala Xaa<sub>19</sub> Xaa<sub>20</sub>  
Xaa<sub>21</sub> Xaa<sub>22</sub> Xaa<sub>23</sub> Xaa<sub>24</sub> Xaa<sub>25</sub> Xaa<sub>26</sub> X<sub>1</sub> -Z<sub>1</sub>; wherein

10

Xaa<sub>1</sub> is His, Arg or Tyr or 4-imidazopropionyl;

Xaa<sub>2</sub> is Ser, Gly, Ala or Thr;

Xaa<sub>3</sub> is Asp or Glu;

Xaa<sub>5</sub> is Ala or Thr;

15 Xaa<sub>6</sub> is Ala, Phe, Tyr or naphthylalanine;

Xaa<sub>7</sub> is Thr or Ser;

Xaa<sub>8</sub> is Ala, Ser or Thr;

Xaa<sub>9</sub> is Asp or Glu;

Xaa<sub>10</sub> is Ala, Leu, Ile, Val, pentylglycine or Met;

20 Xaa<sub>11</sub> is Ala or Ser;

Xaa<sub>12</sub> is Ala or Lys;

Xaa<sub>13</sub> is Ala or Gln;

Xaa<sub>14</sub> is Ala, Leu, Ile, pentylglycine, Val or Met;

Xaa<sub>15</sub> is Ala or Glu;

25 Xaa<sub>16</sub> is Ala or Glu;

Xaa<sub>17</sub> is Ala or Glu;

Xaa<sub>19</sub> is Ala or Val;

Xaa<sub>20</sub> is Ala or Arg;

Xaa<sub>21</sub> is Ala, Leu or Lys-NH<sup>+</sup>-R where R is Lys, Arg, C<sub>1</sub>-C<sub>10</sub>

30 straight chain or branched alkanoyl or cycloalkylalkanoyl;

Xaa<sub>22</sub> is Phe, Tyr or naphthylalanine;

Xaa<sub>23</sub> is Ile, Val, Leu, pentylglycine, tert-butylglycine  
or Met;

Xaa<sub>24</sub> is Ala, Glu or Asp;

Xaa<sub>25</sub> is Ala, Trp, Phe, Tyr or naphthylalanine;

Xaa<sub>26</sub> is Ala or Leu;

5 X<sub>1</sub> is Lys Asn, Asn Lys, Lys-NH<sup>e</sup>-R Asn, Asn Lys-NH<sup>e</sup>-R, Lys-NH<sup>e</sup>-  
R Ala, Ala Lys-NH<sup>e</sup>-R where R is Lys, Arg, C<sub>1</sub>-C<sub>10</sub> straight  
chain or branched alkanoyl or cycloalkylalkanoyl

Z<sub>1</sub> is -OH,

-NH<sub>2</sub>,

Gly-Z<sub>2</sub>,

10 Gly Gly-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub>-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly-Z<sub>2</sub>,

15 Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub>-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub>-Z<sub>2</sub> or

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub>-Z<sub>2</sub>;

wherein

20 Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently  
selected from the group consisting of Pro,  
homoproline, 3Hyp, 4Hyp, thioproline,  
N-alkylglycine, N-alkylpentylglycine and  
N-alkylalanine; and

25 Z<sub>2</sub> is -OH or -NH<sub>2</sub>;

provided that no more than three of Xaa<sub>3</sub>, Xaa<sub>5</sub>, Xaa<sub>6</sub>, Xaa<sub>8</sub>,  
Xaa<sub>10</sub>, Xaa<sub>11</sub>, Xaa<sub>12</sub>, Xaa<sub>13</sub>, Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>, Xaa<sub>17</sub>, Xaa<sub>19</sub>,  
Xaa<sub>20</sub>, Xaa<sub>21</sub>, Xaa<sub>24</sub>, Xaa<sub>25</sub>, and Xaa<sub>26</sub> are Ala. Also within the  
scope of the present invention are pharmaceutically  
30 acceptable salts of the compound of formula (V) and  
pharmaceutical compositions including said compounds and  
salts thereof.

Preferred exendin agonist compounds of formula (V)

include those wherein Xaa<sub>1</sub> is His, Tyr or 4-imidazopropionyl.  
More preferably Xaa<sub>1</sub> is His.

Preferred are those compounds of formula (V) wherein  
Xaa<sub>1</sub> is 4-imidazopropionyl.

5 Preferred are those compounds of formula (V) wherein  
Xaa<sub>2</sub> is Gly.

Preferred compounds of formula (V) are those wherein  
Xaa<sub>14</sub> is Leu, pentylglycine or Met.

Preferred compounds of formula (V) are those wherein  
10 Xaa<sub>25</sub> is Trp or Phe.

According to one aspect, preferred are compounds of  
formula (V) wherein Xaa<sub>6</sub> is Phe or naphthylalanine; and Xaa<sub>22</sub>  
is Phe or naphthylalanine; and Xaa<sub>23</sub> is Ile or Val. More  
preferably, Z<sub>1</sub> is -NH<sub>2</sub>. According to one aspect, especially  
15 preferred are such compounds of formula (V) wherein Xaa<sub>31</sub>,  
Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently selected from the  
group consisting of Pro, homoproline, thioproline and N-  
alkylalanine. More preferably, Z<sub>2</sub> is -NH<sub>2</sub>.

Preferred compounds of formula (V) include those  
20 wherein X<sub>1</sub> is Lys Asn, Lys-NH<sup>c</sup>-R Asn, or Lys-NH<sup>c</sup>-R Ala where R  
is Lys, Arg, C<sub>1</sub>-C<sub>10</sub> straight chain or branched alkanoyl.  
Preferred compounds of formula (V) include compounds  
described in PCT application Serial No. PCT/US98/24210,  
filed November 13, 1998, entitled "Novel Exendin Agonist  
25 Compounds" and identified therein as Compound Nos. 62-69.

Preferred such exendin agonist compounds include those  
wherein Xaa<sub>1</sub> is His, Ala or Norval. More preferably Xaa<sub>1</sub> is  
His or Ala. Most preferably Xaa<sub>1</sub> is His.

Preferred are those compounds of formula (V) wherein  
30 Xaa<sub>2</sub> is Gly.

Preferred are those compounds of formula (V) wherein  
Xaa<sub>3</sub> is Ala.

Preferred are those compounds of formula (V) wherein Xaa<sub>4</sub> is Ala.

Preferred are those compounds of formula (V) wherein Xaa<sub>9</sub> is Ala.

5 Preferred are those compounds of formula (V) wherein Xaa<sub>14</sub> is Leu, pentylglycine or Met.

Preferred compounds of formula (V) are those wherein Xaa<sub>25</sub> is Trp or Phe.

10 Preferred compounds of formula (V) are those where Xaa<sub>6</sub> is Ala, Phe or naphthylalanine; Xaa<sub>22</sub> is Phe or naphthylalanine; and Xaa<sub>23</sub> is Ile or Val.

Preferred are compounds of formula (V) wherein Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently selected from Pro, homoproline, thioproline and N-alkylalanine.

15 Preferably Z<sub>1</sub> is -NH<sub>2</sub>.

Preferably Z<sub>2</sub> is -NH<sub>2</sub>.

According to one aspect, preferred are compounds of formula (V) wherein Xaa<sub>1</sub> is Ala, His or Tyr, more preferably Ala or His; Xaa<sub>2</sub> is Ala or Gly; Xaa<sub>6</sub> is Phe or naphthylalanine; Xaa<sub>14</sub> is Ala, Leu, pentylglycine or Met; Xaa<sub>22</sub> is Phe or naphthylalanine; Xaa<sub>23</sub> is Ile or Val; Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently selected from Pro, homoproline, thioproline or N-alkylalanine; and Xaa<sub>39</sub> is Ser or Tyr, more preferably Ser. More preferably Z<sub>1</sub> is -NH<sub>2</sub>.

25 According to an especially preferred aspect, especially preferred compounds include those of formula (V) wherein: Xaa<sub>1</sub> is His or Ala; Xaa<sub>2</sub> is Gly or Ala; Xaa<sub>3</sub> is Ala, Asp or Glu; Xaa<sub>4</sub> is Ala or Gly; Xaa<sub>5</sub> is Ala or Thr; Xaa<sub>6</sub> is Phe or naphthylalanine; Xaa<sub>7</sub> is Thr or Ser; Xaa<sub>8</sub> is Ala, Ser or Thr; Xaa<sub>9</sub> is Ala, Asp or Glu; Xaa<sub>10</sub> is Ala, Leu or pentylglycine; Xaa<sub>11</sub> is Ala or Ser; Xaa<sub>12</sub> is Ala or Lys; Xaa<sub>13</sub> is Ala or Gln; Xaa<sub>14</sub> is Ala, Leu, Met or pentylglycine; Xaa<sub>15</sub> is Ala or Glu; Xaa<sub>16</sub> is Ala or Glu; Xaa<sub>17</sub> is Ala or Glu; Xaa<sub>19</sub> is Ala or Val;

Xaa<sub>20</sub> is Ala or Arg; Xaa<sub>21</sub> is Ala or Leu; Xaa<sub>22</sub> is Phe or naphthylalanine; Xaa<sub>23</sub> is Ile, Val or tert-butylglycine; Xaa<sub>24</sub> is Ala, Glu or Asp; Xaa<sub>25</sub> is Ala, Trp or Phe; Xaa<sub>26</sub> is Ala or Leu; Xaa<sub>27</sub> is Ala or Lys; Xaa<sub>28</sub> is Ala or Asn; Z<sub>1</sub> is -OH, -NH<sub>2</sub>, Gly-Z<sub>2</sub>, Gly Gly-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub>-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser Gly-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub>-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub>-Z<sub>2</sub>, Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub>-Z<sub>2</sub> or Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub> Xaa<sub>39</sub>-Z<sub>2</sub>; Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> being independently Pro homoproline, thioproline or N-methylalanine; and Z<sub>2</sub> being -OH or -NH<sub>2</sub>; provided that no more than three of Xaa<sub>3</sub>, Xaa<sub>5</sub>, Xaa<sub>6</sub>, Xaa<sub>8</sub>, Xaa<sub>10</sub>, Xaa<sub>11</sub>, Xaa<sub>12</sub>, Xaa<sub>13</sub>, Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>, Xaa<sub>17</sub>, Xaa<sub>19</sub>, Xaa<sub>20</sub>, Xaa<sub>21</sub>, Xaa<sub>24</sub>, Xaa<sub>25</sub>, Xaa<sub>26</sub>, Xaa<sub>27</sub> and Xaa<sub>28</sub> are Ala; and provided also that, if Xaa<sub>1</sub> is His, Arg or Tyr, then at least one of Xaa<sub>3</sub>, Xaa<sub>4</sub> and Xaa<sub>9</sub> is Ala. Especially preferred compounds of formula (V) include those described in PCT application Serial No. PCT/US98/24210, filed November 13, 1998, entitled "Novel Exendin Agonist Compounds" and having the amino acid sequences identified therein as SEQ. ID. NOS. 5-93.

According to an especially preferred aspect, provided are compounds of formula (V) where Xaa<sub>14</sub> is Ala, Leu, Ile, Val or pentylglycine, more preferably Leu or pentylglycine, and Xaa<sub>25</sub> is Ala, Phe, Tyr or naphthylalanine, more preferably Phe or naphthylalanine. These compounds will be less susceptible to oxidative degradation, both in vitro and in vivo, as well as during synthesis of the compound.

#### FORMULA VI

Also provided are peptide compounds described in PCT Application Serial No. PCT/US98/24273, filed November 13, 1998, entitled "Novel Exendin Agonist Compounds", including



compounds of the formula (VI) [SEQ. ID. NO. 46]:

5

10

Xaa<sub>1</sub> Xaa<sub>2</sub> Xaa<sub>3</sub> Xaa<sub>4</sub> Xaa<sub>5</sub> Xaa<sub>6</sub> Xaa<sub>7</sub> Xaa<sub>8</sub> Xaa<sub>9</sub> Xaa<sub>10</sub>

Xaa<sub>11</sub> Xaa<sub>12</sub> Xaa<sub>13</sub> Xaa<sub>14</sub> Xaa<sub>15</sub> Xaa<sub>16</sub> Xaa<sub>17</sub> Ala Xaa<sub>19</sub> Xaa<sub>20</sub>

5 Xaa<sub>21</sub> Xaa<sub>22</sub> Xaa<sub>23</sub> Xaa<sub>24</sub> Xaa<sub>25</sub> Xaa<sub>26</sub> X<sub>1</sub>-Z<sub>1</sub>; wherein

Xaa<sub>1</sub> is His, Arg, Tyr, Ala, Norval, Val, Norleu or 4-imidazopropionyl;

Xaa<sub>2</sub> is Ser, Gly, Ala or Thr;

Xaa<sub>3</sub> is Ala, Asp or Glu;

10 Xaa<sub>4</sub> is Ala, Norval, Val, Norleu or Gly;

Xaa<sub>5</sub> is Ala or Thr;

Xaa<sub>6</sub> is Phe, Tyr or naphthylalanine;

Xaa<sub>7</sub> is Thr or Ser;

Xaa<sub>8</sub> is Ala, Ser or Thr;

15 Xaa<sub>9</sub> is Ala, Norval, Val, Norleu, Asp or Glu;

Xaa<sub>10</sub> is Ala, Leu, Ile, Val, pentylglycine or Met;

Xaa<sub>11</sub> is Ala or Ser;

Xaa<sub>12</sub> is Ala or Lys;

Xaa<sub>13</sub> is Ala or Gln;

20 Xaa<sub>14</sub> is Ala, Leu, Ile, pentylglycine, Val or Met;

Xaa<sub>15</sub> is Ala or Glu;

Xaa<sub>16</sub> is Ala or Glu;

Xaa<sub>17</sub> is Ala or Glu;

Xaa<sub>19</sub> is Ala or Val;

25 Xaa<sub>20</sub> is Ala or Arg;

Xaa<sub>21</sub> is Ala, Leu or Lys-NH<sup>e</sup>-R where R is Lys, Arg, C<sup>1-10</sup> straight chain or branched alkanoyl or cycloalylel-alkanoyl;

Xaa<sub>22</sub> is Phe, Tyr or naphthylalanine;

Xaa<sub>23</sub> is Ile, Val, Leu, pentylglycine, tert-butylglycine or

30 Met;

Xaa<sub>24</sub> is Ala, Glu or Asp;

Xaa<sub>25</sub> is Ala, Trp, Phe, Tyr or naphthylalanine;

Xaa<sub>26</sub> is Ala or Leu;

X<sub>1</sub> is Lys Asn, Asn Lys, Lys-NH<sup>ε</sup>-R Asn, Asn Lys-NH<sup>ε</sup>-R, Lys-NH<sup>ε</sup>-R Ala, Ala Lys-NH<sup>ε</sup>-R where R is Lys, Arg, C<sub>1</sub>-C<sub>10</sub> straight chain or branched alkanoyl or cycloalkylalkanoyl

Z<sub>1</sub> is -OH,

5 -NH<sub>2</sub>,

Gly-Z<sub>2</sub>,

Gly Gly-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub>-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser-Z<sub>2</sub>,

10 Gly Gly Xaa<sub>31</sub> Ser Ser-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub>-Z<sub>2</sub>,

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub>-Z<sub>2</sub>,

15 Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub>-Z<sub>2</sub> or

Gly Gly Xaa<sub>31</sub> Ser Ser Gly Ala Xaa<sub>36</sub> Xaa<sub>37</sub> Xaa<sub>38</sub> Xaa<sub>39</sub>-Z<sub>2</sub>;

wherein

Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently

selected from the group consisting of Pro,

20 homoproline, 3Hyp, 4Hyp, thioproline,

N-alkylglycine, N-alkylpentylglycine and

N-alkylalanine; and

Z<sub>2</sub> is -OH or -NH<sub>2</sub>;

provided that no more than three of Xaa<sub>3</sub>, Xaa<sub>4</sub>, Xaa<sub>5</sub>, Xaa<sub>6</sub>,

25 Xaa<sub>8</sub>, Xaa<sub>9</sub>, Xaa<sub>10</sub>, Xaa<sub>11</sub>, Xaa<sub>12</sub>, Xaa<sub>13</sub>, Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>,

Xaa<sub>17</sub>, Xaa<sub>19</sub>, Xaa<sub>20</sub>, Xaa<sub>21</sub>, Xaa<sub>24</sub>, Xaa<sub>25</sub>, Xaa<sub>26</sub>, are Ala; and

provided also that, if Xaa<sub>1</sub> is His, Arg, Tyr, or 4-

imidazopropionyl then at least one of Xaa<sub>3</sub>, Xaa<sub>4</sub> and Xaa<sub>9</sub> is Ala.

30 Preferred compounds of formula (VI) include those wherein Xaa<sub>1</sub> is His, Ala, Norval or 4-imidazopropionyl.

Preferably, Xaa<sub>1</sub> is His, or 4-imidazopropionyl or Ala, more preferably His or 4-imidazopropionyl.

Preferred compounds of formula (VI) include those wherein Xaa<sub>2</sub> is Gly.

Preferred compounds of formula (VI) include those wherein Xaa<sub>4</sub> is Ala.

5 Preferred compounds of formula (VI) include those wherein Xaa<sub>9</sub> is Ala.

Preferred compounds of formula (VI) include those wherein Xaa<sub>14</sub> is Leu, pentylglycine or Met.

10 Preferred compounds of formula (VI) include those wherein Xaa<sub>25</sub> is Trp or Phe.

Preferred compounds of formula (VI) include those wherein Xaa<sub>6</sub> is Ala, Phe or naphthylalanine; Xaa<sub>22</sub> is Phe or naphthylalanine; and Xaa<sub>23</sub> is Ile or Val.

15 Preferred compounds of formula (VI) include those wherein Z<sub>1</sub> is -NH<sub>2</sub>.

Preferred compounds of formula (VI) include those wherein Xaa<sub>31</sub>, Xaa<sub>36</sub>, Xaa<sub>37</sub> and Xaa<sub>38</sub> are independently selected from the group consisting of Pro, homoproline, thioproline and N-alkylalanine.

20 Preferred compounds of formula (VI) include those wherein Xaa<sub>39</sub> is Ser or Tyr, preferably Ser.

Preferred compounds of formula (VI) include those wherein Z<sub>2</sub> is -NH<sub>2</sub>.

25 Preferred compounds of formula (VI) include those wherein Z<sub>1</sub> is -NH<sub>2</sub>.

Preferred compounds of formula (VI) include those wherein Xaa<sub>21</sub> is Lys-NH<sup>e</sup>-R where R is Lys, Arg, C<sub>1</sub>-C<sub>10</sub> straight chain or branched alkanoyl.

30 Preferred compounds of formula (VI) include those wherein X<sub>1</sub> is Lys Asn, Lys-NH<sup>e</sup>-R Asn, or Lys-NH<sup>e</sup>-R Ala where R is Lys, Arg, C<sub>1</sub>-C<sub>10</sub> straight chain or branched alkanoyl.

Preferred compounds of formula (VI) include those described in PCT Application Serial No. PCT/US98/24273,

filed November 13, 1998, entitled "Novel Exendin Agonist Compounds" as having an amino acid sequence selected from those identified therein as SEQ. ID. NOS. 95-110.

# 5 FORMULA VII

Compounds particularly useful according to the present invention are exendin agonist compounds described in U.S. Patent Application Serial No. 09/003,869, filed January 7, 1998, entitled "Use of Exendins And Agonists Thereof For The

10 Reduction of Food Intake", including compounds of the formula (VII) [SEQ. ID. NO. 47]:

1	5	10
Xaa <sub>1</sub>	Xaa <sub>2</sub>	Xaa <sub>3</sub>
Gly	Thr	Xaa <sub>4</sub>
Xaa <sub>5</sub>	Xaa <sub>6</sub>	Xaa <sub>7</sub>
Xaa <sub>8</sub>		
15	20	
Ser	Lys	Gln
Xaa <sub>9</sub>	Glu	Glu
Glu	Ala	Val
Arg	Leu	
25	30	
Xaa <sub>10</sub>	Xaa <sub>11</sub>	Xaa <sub>12</sub>
Xaa <sub>13</sub>	Leu	Lys
Asn	Gly	Gly
Xaa <sub>14</sub>		
35		
Ser	Ser	Gly
Ala	Xaa <sub>15</sub>	Xaa <sub>16</sub>
Xaa <sub>17</sub>	Xaa <sub>18</sub>	-Z

20 wherein Xaa<sub>1</sub> is His, Arg or Tyr; Xaa<sub>2</sub> is Ser, Gly, Ala or Thr; Xaa<sub>3</sub> is Asp or Glu; Xaa<sub>4</sub> is Phe, Tyr or naphthalanine; Xaa<sub>5</sub> is Thr or Ser; Xaa<sub>6</sub> is Ser or Thr; Xaa<sub>7</sub> is Asp or Glu; Xaa<sub>8</sub> is Leu, Ile, Val, pentylglycine or Met; Xaa<sub>9</sub> is Leu, Ile, pentylglycine, Val or Met; Xaa<sub>10</sub> is Phe, Tyr or naphthalanine; Xaa<sub>11</sub> is Ile, Val, Leu, pentylglycine, tert-butylglycine or Met; Xaa<sub>12</sub> is Glu or Asp; Xaa<sub>13</sub> is Trp, Phe, Tyr, or naphthylalanine; Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub> and Xaa<sub>17</sub> are independently Pro, homoproline, 3Hyp, 4Hyp, thioproline, N-alkylglycine, N-alkylpentylglycine or N-alkylalanine; Xaa<sub>18</sub>

25 is Ser, Thr or Tyr; and Z is -OH or -NH<sub>2</sub>; with the proviso that the compound does not have the formula of either SEQ. ID. NOS. 1 or 2. Preferred N-alkyl groups for N-

30

alkylglycine, N-alkylpentylglycine and N-alkylalanine include lower alkyl groups preferably of 1 to about 6 carbon atoms, more preferably of 1 to 4 carbon atoms. Suitable compounds include those having amino acid sequences of SEQ. ID. NOS. 10 to 40. Also useful in the present invention are pharmaceutically acceptable salts of the compounds of formula (VII).

Preferred exendin agonist compounds include those wherein Xaa<sub>1</sub> is His or Tyr. More preferably Xaa<sub>1</sub> is His.

Preferred are those compounds wherein Xaa<sub>2</sub> is Gly.

Preferred are those compounds wherein Xaa<sub>9</sub> is Leu, pentylglycine or Met.

Preferred compounds include those wherein Xaa<sub>13</sub> is Trp or Phe.

Also preferred are compounds where Xaa<sub>4</sub> is Phe or naphthalanine; Xaa<sub>11</sub> is Ile or Val and Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub> and Xaa<sub>17</sub> are independently selected from Pro, homoproline, thioproline or N-alkylalanine. Preferably N-alkylalanine has a N-alkyl group of 1 to about 6 carbon atoms.

According to an especially preferred aspect, Xaa<sub>15</sub>, Xaa<sub>16</sub> and Xaa<sub>17</sub> are the same amino acid residue.

Preferred are compounds wherein Xaa<sub>18</sub> is Ser or Tyr, more preferably Ser.

Preferably Z is -NH<sub>2</sub>.

According to one aspect, preferred are compounds of formula (VII) wherein Xaa<sub>1</sub> is His or Tyr, more preferably His; Xaa<sub>2</sub> is Gly; Xaa<sub>4</sub> is Phe or naphthalanine; Xaa<sub>9</sub> is Leu, pentylglycine or Met; Xaa<sub>10</sub> is Phe or naphthalanine; Xaa<sub>11</sub> is Ile or Val; Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub> and Xaa<sub>17</sub> are independently selected from Pro, homoproline, thioproline or N-alkylalanine; and Xaa<sub>18</sub> is Ser or Tyr, more preferably Ser. More preferably Z is -NH<sub>2</sub>.

According to an especially preferred aspect, especially preferred compounds include those of formula (VII) wherein: Xaa<sub>1</sub> is His or Arg; Xaa<sub>2</sub> is Gly; Xaa<sub>3</sub> is Asp or Glu; Xaa<sub>4</sub> is Phe or naphthylalanine; Xaa<sub>5</sub> is Thr or Ser; Xaa<sub>6</sub> is Ser or Thr; Xaa<sub>7</sub> is Asp or Glu; Xaa<sub>8</sub> is Leu or pentylglycine; Xaa<sub>9</sub> is Leu or pentylglycine; Xaa<sub>10</sub> is Phe or naphthylalanine; Xaa<sub>11</sub> is Ile, Val or t-butyltylglycine; Xaa<sub>12</sub> is Glu or Asp; Xaa<sub>13</sub> is Trp or Phe; Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub>, and Xaa<sub>17</sub> are independently Pro, homoproline, thioproline, or N-methylalanine; Xaa<sub>18</sub> is Ser or Tyr; and Z is -OH or -NH<sub>2</sub>; with the proviso that the compound does not have the formula of either SEQ. ID. NOS. 1 or 2. More preferably Z is -NH<sub>2</sub>. Especially preferred compounds include those having the amino acid sequence of SEQ. ID. NOS. 10, 11, 22, 23, 24, 27, 29, 36, 37 and 40.

According to an especially preferred aspect, provided are compounds where Xaa<sub>9</sub> is Leu, Ile, Val or pentylglycine, more preferably Leu or pentylglycine, and Xaa<sub>13</sub> is Phe, Tyr or naphthylalanine, more preferably Phe or naphthylalanine. These compounds are believed to exhibit advantageous duration of action and to be less subject to oxidative degradation, both in vitro and in vivo, as well as during synthesis of the compound.

#### 25 FORMULA VIII

Also provided are compounds described in PCT Application Serial No. PCT/US98/16387, filed August 6, 1998, entitled "Novel Exendin Agonist Compounds", including compounds of the formula (VIII) [SEQ. ID. NO. 48]:

30	1	5	10
	Xaa <sub>1</sub>	Xaa <sub>2</sub>	Xaa <sub>3</sub>
	Gly	Thr	Xaa <sub>4</sub>
	Xaa <sub>5</sub>	Xaa <sub>6</sub>	Xaa <sub>7</sub>
	Xaa <sub>8</sub>		
	15	20	
	Ser	Lys	Gln
	Xaa <sub>9</sub>	Glu	Glu
	Glu	Ala	Val
	Arg	Leu	

25

30

Xaa<sub>10</sub> Xaa<sub>11</sub> Xaa<sub>12</sub> Xaa<sub>13</sub> Leu X<sub>1</sub> Gly Gly Xaa<sub>14</sub>

35

Ser Ser Gly Ala Xaa<sub>15</sub> Xaa<sub>16</sub> Xaa<sub>17</sub> Xaa<sub>18</sub>-Z

- 5 wherein Xaa<sub>1</sub> is His, Arg, Tyr or 4-imidazopropionyl; Xaa<sub>2</sub> is Ser, Gly, Ala or Thr; Xaa<sub>3</sub> is Asp or Glu; Xaa<sub>4</sub> is Phe, Tyr or naphthylalanine; Xaa<sub>5</sub> is Thr or Ser; Xaa<sub>6</sub> is Ser or Thr; Xaa<sub>7</sub> is Asp or Glu; Xaa<sub>8</sub> is Leu, Ile, Val, pentylglycine or Met; Xaa<sub>9</sub> is Leu, Ile, pentylglycine, Val or Met; Xaa<sub>10</sub> is Phe,
- 10 Tyr or naphthylalanine; Xaa<sub>11</sub> is Ile, Val, Leu, pentylglycine, tert-butylglycine or Met; Xaa<sub>12</sub> is Glu or Asp; Xaa<sub>13</sub> is Trp, Phe, Tyr, or naphthylalanine; X<sub>1</sub> is Lys Asn, Asn Lys, Lys-NH<sup>e</sup>-R Asn, Asn Lys-NH<sup>e</sup>-R where R is Lys, Arg, C<sub>1</sub>-C<sub>10</sub> straight chain or branched alkanoyl or
- 15 cycloalkylalkanoyl; Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub> and Xaa<sub>17</sub> are independently Pro, homoproline, 3Hyp, 4Hyp, thioproline, N-alkylglycine, N-alkylpentylglycine or N-alkylalanine; Xaa<sub>18</sub> is Ser, Thr or Tyr; and Z is -OH or -NH<sub>2</sub>; with the proviso that the compound does not have the formula of either SEQ.
- 20 ID. NOS. 1 or 2. Suitable compounds of formula (VIII) include compounds described in PCT Application Serial No. PCT/US98/16387, filed August 6, 1998, entitled "Novel Exendin Agonist Compounds" having the amino acid sequences of SEQ. ID. NOS. 37-40 therein.

- 25 Preferred exendin agonist compounds of formula (VIII) include those wherein Xaa<sub>1</sub> is His, Tyr or 4-imidazopropionyl. More preferably, Xaa<sub>1</sub> is His or 4-imidazopropionyl.

Preferred are those compounds of formula (VIII) wherein Xaa<sub>2</sub> is Gly.

- 30 Preferred are those compounds of formula (VIII) wherein Xaa<sub>9</sub> is Leu, pentylglycine or Met.

Preferred are those compounds of formula (VIII) wherein Xaa<sub>13</sub> is Trp or Phe.

Preferred are those compounds of formula (VIII) wherein  $X_1$  is Lys Asn, or Lys-NH<sup>e</sup>-R Asn, where R is Lys, Arg, C<sub>1</sub>-C<sub>10</sub> straight chain or branched alkanoyl.

Also preferred are compounds of formula (VIII) wherein  
5 Xaa<sub>4</sub> is Phe or naphthylalanine; Xaa<sub>10</sub> is Phe or naphthylalanine; Xaa<sub>11</sub> is Ile or Val and Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub> and Xaa<sub>17</sub> are independently selected from Pro, homoproline, thioproline or N-alkylalanine. According to an especially preferred aspect, Xaa<sub>18</sub> is Ser or Tyr. Preferred are those  
10 such compounds wherein Xaa<sub>18</sub> is Ser. Preferably, Z is -NH<sub>2</sub>.

According to one preferred aspect, preferred are compounds of formula (VIII) wherein Xaa<sub>4</sub> is Phe or naphthylalanine; Xaa<sub>10</sub> is Phe or naphthylalanine; Xaa<sub>11</sub> is Ile or Val,  $X_1$  is Lys Asn, or Lys-NH<sup>e</sup>-R Asn, where R is Lys, Arg,  
15 C<sub>1</sub>-C<sub>10</sub> straight chain or branched alkanoyl and Xaa<sub>14</sub>, Xaa<sub>15</sub>, Xaa<sub>16</sub> and Xaa<sub>17</sub> are independently selected from Pro, homoproline, thioproline or N-alkylalanine.

#### Preparation of Modified Exendins And Exendin Agonists

20 The modified exendins and exendin agonists of the present invention may be made by linking one or more polyethylene glycol polymers to an exendin or exendin agonist. The synthesis of exendins and exendin agonists is thus described first, followed by methodology for linking  
25 the polyethylene glycol polymer(s) to the exendin or exendin agonist.

#### Preparation of Exendins And Exendin Agonists

30 Exendins and exendin agonist compounds such as exendin analogs and exendin derivatives, described herein may be prepared through peptide purification as described in, for example, Eng, et al., J. Biol. Chem. 265:20259-62, 1990; and Eng, et al., J. Biol. Chem. 267:7402-05, 1992, hereby



incorporated by reference herein. Alternatively, exendins and exendin agonist peptides may be prepared by methods known to those skilled in the art, for example, as described in Raufman, et al. (J. Biol. Chem. 267:21432-37, 1992),

5 hereby incorporated by reference herein, using standard solid-phase peptide synthesis techniques and preferably an automated or semiautomated peptide synthesizer. The compounds that constitute active ingredients of the formulations and dosages of the present invention may be prepared using standard solid-phase peptide synthesis techniques and preferably an automated or semiautomated peptide synthesizer. Typically, using such techniques, an  $\alpha$ -N-carbamoyl protected amino acid and an amino acid attached to the growing peptide chain on a resin are coupled at room temperature in an inert solvent such as dimethylformamide, N-methylpyrrolidinone or methylene chloride in the presence of coupling agents such as dicyclohexylcarbodiimide and 1-hydroxybenzotriazole in the presence of a base such as diisopropylethylamine. The  $\alpha$ -N-carbamoyl protecting group is removed from the resulting peptide-resin using a reagent such as trifluoroacetic acid or piperidine, and the coupling reaction repeated with the next desired N-protected amino acid to be added to the peptide chain. Suitable N-protecting groups are well known in the art, with t-butyloxycarbonyl (tBoc) and fluorenylmethoxycarbonyl (Fmoc) being preferred herein.

The solvents, amino acid derivatives and 4-methylbenzhydryl-amine resin used in the peptide synthesizer may be purchased from Applied Biosystems Inc. (Foster City, CA). The following side-chain protected amino acids may be purchased from Applied Biosystems, Inc.: BSD-112344.1-Arg(Pmc), Boc-Thr(Bzl), Fmoc-Thr(t-Bu), Boc-Ser(Bzl), Fmoc-Ser(t-Bu), Boc-Tyr(BrZ), Fmoc-Tyr(t-Bu), Boc-Lys(Cl-Z),

Fmoc-Lys(Boc), Boc-Glu(Bzl), Fmoc-Glu(t-Bu), Fmoc-His(Trt), Fmoc-Asn(Trt), and Fmoc-Gln(Trt). Boc-His(BOM) may be purchased from Applied Biosystems, Inc. or Bachem Inc.

(Torrance, CA). Anisole, dimethylsulfide, phenol, ethanedithiol, and thioanisole may be obtained from Aldrich Chemical Company (Milwaukee, WI). Air Products and Chemicals (Allentown, PA) supplies HF. Ethyl ether, acetic acid and methanol may be purchased from Fisher Scientific (Pittsburgh, PA).

Solid phase peptide synthesis may be carried out with an automatic peptide synthesizer (Model 430A, Applied Biosystems Inc., Foster City, CA) using the NMP/HOBt (Option 1) system and tBoc or Fmoc chemistry (see, Applied Biosystems User's Manual for the ABI 430A Peptide Synthesizer, Version 1.3B July 1, 1988, section 6, pp. 49-70, Applied Biosystems, Inc., Foster City, CA) with capping. Boc-peptide-resins may be cleaved with HF (-50°C to 0°C, 1 hour). The peptide may be extracted from the resin with alternating water and acetic acid, and the filtrates lyophilized. The Fmoc-peptide resins may be cleaved according to standard methods (Introduction to Cleavage Techniques, Applied Biosystems, Inc., 1990, pp. 6-12). Peptides may also be assembled using an Advanced Chem Tech Synthesizer (Model MPS 350, Louisville, Kentucky).

Peptides may be purified by RP-HPLC (preparative and analytical) using a Waters Delta Prep 3000 system. A C4, C8 or C18 preparative column (10  $\mu$ , 2.2 x 25 cm; Vydac, Hesperia, CA) may be used to isolate peptides, and purity may be determined using a C4, C8 or C18 analytical column (5  $\mu$ , 0.46 x 25 cm; Vydac). Solvents (A=0.1% TFA/water and B=0.1% TFA/CH<sub>3</sub>CN) may be delivered to the analytical column at a flowrate of 1.0 ml/min and to the preparative column at 15 ml/min. Amino acid analyses may be performed on the

Waters Pico Tag system and processed using the Maxima program. Peptides may be hydrolyzed by vapor-phase acid hydrolysis (115°C, 20-24 h). Hydrolysates may be derivatized and analyzed by standard methods (Cohen, et al., The Pico

5 Tag Method: A Manual of Advanced Techniques for Amino Acid Analysis, pp. 11-52, Millipore Corporation, Milford, MA (1989)). Fast atom bombardment analysis may be carried out by M-Scan, Incorporated (West Chester, PA). Mass calibration may be performed using cesium iodide or cesium  
10 iodide/glycerol. Plasma desorption ionization analysis using time of flight detection may be carried out on an Applied Biosystems Bio-Ion 20 mass spectrometer. Electrospray mass spectroscopy may be carried and on a VG-Trio machine.

15 Peptide active ingredient compounds useful in the formulations and dosages of the invention may also be prepared using recombinant DNA techniques, using methods now known in the art. See, e.g., Sambrook et al., Molecular Cloning: A Laboratory Manual, 2d Ed., Cold Spring Harbor  
20 (1989). Alternatively, such compounds may be prepared by homogeneous phase peptide synthesis methods. Non-peptide compounds useful in the present invention may be prepared by art-known methods. For example, phosphate-containing amino acids and peptides containing such amino acids, may be  
25 prepared using methods known in the art. See, e.g., Bartlett and Landen, Biorg. Chem. 14:356-377 (1986).

#### Conjugation of Polyethylene Glycol Polymers

There are several strategies for coupling PEG to  
30 peptides/proteins. See, Int. J. Hematology 68:1 (1998); Bioconjugate Chem. 6:150 (1995); and Crit. Rev. Therap. Drug Carrier Sys. 9:249 (1992) all of which are incorporated herein by reference in their entirety. Those skilled in the

art, therefore, will be able to utilize such well-known techniques for linking one or more polyethylene glycol polymers to the exendins and exendin agonists described herein. Suitable polyethylene glycol polymers typically are commercially available or may be made by techniques well known to those skilled in the art. The polyethylene glycol polymers preferably have molecular weights between 500 and 20,000 and may be branched or straight chain polymers.

The attachment of a PEG on an intact peptide or protein can be accomplished by coupling to amino, carboxyl or thiol groups. These groups will typically be the N and C termini and on the side chains of such naturally occurring amino acids as lysine, aspartic acid, glutamic acid and cysteine. Since exendin 4 and other exendins and exendin agonists can be prepared by solid phase peptide chemistry techniques, a variety of moieties containing diamino and dicarboxylic groups with orthogonal protecting groups can be introduced for conjugation to PEG.

The present invention also provides for conjugation of an exendin or exendin agonist to one or more polymers other than polyethylene glycol which can regulate kidney clearance in a manner similar to polyethylene glycol. Examples of such polymers include albumin and gelatin. See, Gombotz and Pettit, *Bioconjugate Chem.*, 6:332-351, 1995, which is incorporated herein by reference in its entirety.

#### Utility

The formulations and dosages described herein are useful in view of their pharmacological properties. In particular, the compounds described herein possess activity as agents to reduce glucagon levels and suppress glucagon secretion, as evidenced by the ability to lower glucagon levels in animals and humans. They can be used to treat

conditions or diseases that can be alleviated by reducing glucagon levels and suppressing glucagon secretion.

The compounds referenced above may form salts with various inorganic and organic acids and bases. Such salts include salts prepared with organic and inorganic acids, for example, HCl, HBr, H<sub>2</sub>SO<sub>4</sub>, H<sub>3</sub>PO<sub>4</sub>, trifluoroacetic acid, acetic acid, formic acid, methanesulfonic acid, toluenesulfonic acid, maleic acid, fumaric acid and camphorsulfonic acid. Salts prepared with bases include ammonium salts, alkali metal salts, e.g., sodium and potassium salts, and alkali earth salts, e.g., calcium and magnesium salts. Acetate, hydrochloride, and trifluoroacetate salts are preferred. The salts may be formed by conventional means, as by reacting the free acid or base forms of the product with one or more equivalents of the appropriate base or acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water which is then removed in vacuo or by freeze-drying or by exchanging the ions of an existing salt for another ion on a suitable ion exchange resin.

#### Formulation and Administration

Modified exendin and exendin agonist formulations and dosages of the invention are useful in view of their exendin-like effects, and may conveniently be provided in the form of formulations suitable for parenteral (including intravenous, intramuscular and subcutaneous) administration. Also described herein are formulations and dosages useful in alternative delivery routes, including oral, nasal, buccal, sublingual and pulmonary.

The feasibility of alternate routes of delivery for exendin-4 has been explored by measuring exendin-4 in the circulation in conjunction with observation of a biologic response, such as plasma glucose lowering in diabetic

animals, after administration. Passage of exendin-4 has been investigated across several surfaces, the respiratory tract (nasal, tracheal and pulmonary routes) and the gut (sublingual, gavage and intraduodenal routes). Biologic effect and appearance of exendin-4 in blood have been observed with each route of administration via the respiratory tract, and with sublingual and gavaged peptide via the gastrointestinal tract. Intra-tracheal administration, nasal administration, administration via the gut, and sublingual administration have all been described.

In some cases, it will be convenient to provide a modified exendin or exendin agonist and another anti-glucagon agent, such as an amylin or an amylin agonist, in a single composition or solution for administration together.

In other cases, it may be more advantageous to administer another anti-glucagon agent separately from the exendin, exendin agonist, or modified exendin or exendin agonist. In yet other cases, it may be beneficial to provide an exendin, exendin agonist, or modified exendin or exendin agonist either co-formulated or separately with other glucagon lowering agents such as amylin. A suitable administration format may best be determined by a medical practitioner for each patient individually. Suitable pharmaceutically acceptable carriers and their formulation are described in standard formulation treatises, e.g., Remington's Pharmaceutical Sciences by E.W. Martin. See also Wang, Y.J. and Hanson, M.A. "Parenteral Formulations of Proteins and Peptides: Stability and Stabilizers," Journal of Parenteral Science and Technology, Technical Report No. 10, Supp. 42:2S (1988).

Compounds useful in the invention can be provided as parenteral compositions for injection or infusion. They can, for example, be suspended in an inert oil, suitably a

vegetable oil such as sesame, peanut, olive oil, or other acceptable carrier. Preferably, they are suspended in an aqueous carrier, for example, in an isotonic buffer solution at a pH of about 5.6 to 7.4. These compositions may be  
5 sterilized by conventional sterilization techniques, or may be sterile filtered. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH buffering agents. Useful buffers include for example,  
10 sodium acetate/acetic acid buffers. A form of repository or "depot" slow release preparation may be used so that therapeutically effective amounts of the preparation are delivered into the bloodstream over many hours or days following transdermal injection or delivery.

15 The desired isotonicity may be accomplished using sodium chloride or other pharmaceutically acceptable agents such as dextrose, boric acid, sodium tartrate, propylene glycol, polyols (such as mannitol and sorbitol), or other inorganic or organic solutes. Sodium chloride is preferred  
20 particularly for buffers containing sodium ions.

The claimed compounds can also be formulated as pharmaceutically acceptable salts (e.g., acid addition salts) and/or complexes thereof. Pharmaceutically acceptable salts are non-toxic salts at the concentration at  
25 which they are administered. The preparation of such salts can facilitate the pharmacological use by altering the physical-chemical characteristics of the composition without preventing the composition from exerting its physiological effect. Examples of useful alterations in physical  
30 properties include lowering the melting point to facilitate transmucosal administration and increasing the solubility to facilitate the administration of higher concentrations of the drug.

Pharmaceutically acceptable salts include acid addition salts such as those containing sulfate, hydrochloride, phosphate, sulfamate, acetate, citrate, lactate, tartrate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate, cyclohexylsulfamate and quinate.

Pharmaceutically acceptable salts can be obtained from acids such as hydrochloric acid, sulfuric acid, phosphoric acid, sulfamic acid, acetic acid, citric acid, lactic acid, tartaric acid, malonic acid, methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, *p*-toluenesulfonic acid, cyclohexylsulfamic acid, and quinic acid. Such salts may be prepared by, for example, reacting the free acid or base forms of the product with one or more equivalents of the appropriate base or acid in a solvent or medium in which the salt is insoluble, or in a solvent such as water which is then removed in vacuo or by freeze-drying or by exchanging the ions of an existing salt for another ion on a suitable ion exchange resin.

Carriers or excipients can also be used to facilitate administration of the compound. Examples of carriers and excipients include calcium carbonate, calcium phosphate, various sugars such as lactose, glucose, or sucrose, or types of starch, cellulose derivatives, gelatin, vegetable oils, polyethylene glycols and physiologically compatible solvents. The compositions or pharmaceutical composition can be administered by different routes including intravenously, intraperitoneal, subcutaneous, and intramuscular, orally, topically, or transmucosally.

If desired, solutions of the above compositions may be thickened with a thickening agent such as methylcellulose. They may be prepared in emulsified form, either water in oil or oil in water. Any of a wide variety of pharmaceutically acceptable emulsifying agents may be employed including, for



example, acacia powder, a non-ionic surfactant (such as a Tween), or an ionic surfactant (such as alkali polyether alcohol sulfates or sulfonates, e.g., a Triton).

Compositions useful in the invention are prepared by  
5 mixing the ingredients following generally accepted procedures. For example, the selected components may be simply mixed in a blender or other standard device to produce a concentrated mixture which may then be adjusted to the final concentration and viscosity by the addition of  
10 water or thickening agent and possibly a buffer to control pH or an additional solute to control tonicity.

For use by the physician, the compounds will be provided in dosage unit form containing an amount of an exendin, exendin agonist, or modified exendin or exendin  
15 agonist, with or without another anti-glucagon agent. Therapeutically effective amounts of an exendin, exendin agonist, or modified exendin or exendin agonist for use in the control of glucagon and in conditions in which glucagon levels are beneficially lowered or regulated are those that  
20 decrease post-prandial blood glucagon levels as desired. In diabetic or glucose intolerant individuals, plasma glucagon levels may be higher than in normal individuals. In such individuals, beneficial reduction or "smoothing" of post-prandial blood glucagon levels, may be obtained. As will be  
25 recognized by those in the field, an effective amount of therapeutic agent will vary with many factors including the age and weight of the patient, the patient's physical condition, the glucagon level or level of inhibition of glucagon suppression to be obtained, and other factors.

30 Such pharmaceutical compositions are useful in causing glucagon to be lowered in a subject and may be used as well in other disorders where lowered or suppressed glucagon is beneficially reduced.

The effective daily anti-glucagon dose of the compounds will typically be in the range of 0.01 or 0.03 to about 5 mg/day, preferably about 0.01 or 0.5 to 2 mg/day and more preferably about 0.01 or 0.1 to 1 mg/day, for a 70 kg

5 patient, administered in a single or divided doses. The exact dose to be administered is determined by the attending clinician and is dependent upon where the particular compound lies within the above quoted range, as well as upon the age, weight and condition of the individual.

- 10 Administration should begin at the first sign of symptoms or shortly after diagnosis of, for example, diabetes mellitus as manifested by elevated glucagon. Administration may be by injection, preferably subcutaneous or intramuscular. Orally active compounds may be taken orally, however dosages
- 15 should be increased 5-10 fold.

- Generally, in treating or preventing elevated, inappropriate, or undesired post-prandial blood glucagon levels, the compounds of this invention may be administered to patients in need of such treatment in a dosage ranges
- 20 similar to those given above, however, the compounds are administered more frequently, for example, one, two, or three times a day. Particularly preferred are the exendin and exendin agonist formulations and dosages and routes of administration thereof described commonly owned U.S.
- 25 Provisional Application 60/116,380, entitled "Novel Exendin Agonist Formulations And Methods Of Administration Thereof," filed January 14, 1999 (and the corresponding PCT application claiming priority from it that was filed on January 14, 2000, Serial No. [not yet assigned]), and U.S.
- 30 Provisional Application 60/[not yet assigned], entitled "Use of Exendins and Agonists Thereof for Modulation of Triglyceride Levels and Treatment of Dyslipidemia," filed January 14, 1999, from which this application claims

priority and the disclosures of which have been incorporated by referenced in their entirety as if fully set forth herein.

The optimal formulation and mode of administration of compounds of the present application to a patient depend on factors known in the art such as the particular disease or disorder, the desired effect, and the type of patient. While the compounds will typically be used to treat human patients, they may also be used to treat similar or identical diseases in other vertebrates such as other primates, farm animals such as swine, cattle and poultry, and sports animals and pets such as horses, dogs and cats.

To assist in understanding the present invention the following Examples are included which describe the results of a series of experiments. The experiments relating to this invention should not, of course, be construed as specifically limiting the invention and such variations of the invention, now known or later developed, which would be within the purview of one skilled in the art are considered to fall within the scope of the invention as described herein and hereinafter claimed.

#### EXAMPLE 1 - PREPARATION OF EXENDIN-3

His Ser Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu  
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly  
Pro Ser Ser Gly Ala Pro Pro Pro Ser-NH<sub>2</sub> [SEQ. ID. NO. 1]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.). In general, single-coupling cycles were used throughout the synthesis and Fast Moc (HBTU activation) chemistry was employed. Deprotection (Fmoc group removal) of the growing

peptide chain was achieved using piperidine. Final deprotection of the completed peptide resin was achieved using a mixture of triethylsilane (0.2 mL), ethanedithiol (0.2 mL), anisole (0.2 mL), water (0.2 mL) and

5 trifluoroacetic acid (15 mL) according to standard methods (Introduction to Cleavage Techniques, Applied Biosystems, Inc.) The peptide was precipitated in ether/water (50 mL) and centrifuged. The precipitate was reconstituted in glacial acetic acid and lyophilized. The lyophilized

10 peptide was dissolved in water). Crude purity was about 75%.

Used in purification steps and analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN).

The solution containing peptide was applied to a

15 preparative C-18 column and purified (10% to 40% Solvent B in Solvent A over 40 minutes). Purity of fractions was determined isocratically using a C-18 analytical column. Pure fractions were pooled furnishing the above-identified peptide. Analytical RP-HPLC (gradient 30% to 60% Solvent B

20 in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 19.2 minutes.

#### EXAMPLE 2 - PREPARATION OF EXENDIN-4

25 His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu  
Glu Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Gly  
Pro Ser Ser Gly Ala Pro Pro Pro Ser-NH<sub>2</sub> [SEQ. ID. NO. 2]

The above amidated peptide was assembled on 4-(2'-4'-dimethoxyphenyl)-Fmoc aminomethyl phenoxy acetamide

30 norleucine MBHA resin (Novabiochem, 0.55 mmole/g) using Fmoc-protected amino acids (Applied Biosystems, Inc.), cleaved from the resin, deprotected and purified in a similar way to Exendin-3 as describe in Example 1. Used in

analysis were Solvent A (0.1% TFA in water) and Solvent B (0.1% TFA in ACN). Analytical RP-HPLC (gradient 36% to 46% Solvent B in Solvent A over 30 minutes) of the lyophilized peptide gave product peptide having an observed retention time of 14.9 minutes. Electrospray Mass Spectrometry (M): calculated 4186.6; found 4186.0 to 4186.8 (four lots).

### EXAMPLE 3: CLEARANCE BY THE KIDNEY

The kidney can play a major role in the elimination of some molecules (drugs, peptides, proteins). For some molecules, this process begins when the kidney filters the blood at the glomerulus to produce the ultrafiltrate described below. The glomerular filter discriminates not only on the basis of molecular weight but also by acting as a negatively charged selective barrier, promoting retention of anionic compounds. The free fraction of molecules in the plasma (not protein bound) with a molecular weight less than 5kD and an effective radii less than 15 Å are easily filtered. For larger molecular weight molecules they are filtered on a more restrictive and limited basis, principally by molecular size, structure and net charge. The cutoff point for glomerular filtration lies between albumin (67kD) which is retained and hemoglobin (68kD) which is filtered. Albumin, with an effective radius of about 36 Å is filtered less than 1% at the glomerulus.

Once in the glomerulus a molecule travels to the proximal tubule where it is either reabsorbed or it passes on through the loop of Henle to the distal tubule where collecting ducts drain the filtrate into the bladder. Filtered proteins and peptides are typically cleaved by brush border enzymes in the proximal tubule, from where they are efficiently retrieved by sodium/amino cotransporters (scavenging pumps). Otherwise, molecules which are polar,

ionized and of large molecular weight will not be reabsorbed. Throughout this process metabolizing enzymes in the renal cortex (proximal tubules) may also degrade the molecule into more polar molecules, thereby increasing the probability for excretion into the urine. Many peptide hormones (for example, amylin, calcitonins) are degraded by passage through the renal circulation, presumably by vascular ectoenzymes accessible to the plasma, independently of the process of glomerular filtration. In those examples, rates of peptide clearance from the plasma are similar to the rate of renal plasma flow, which is ~3-fold greater than the rate of glomerular filtration.

Studies performed to identify plasma circulating metabolites of exendin-4 yielded very little evidence of proteolytic degradation; following large intravenous doses in animals, HPLC analysis of plasma showed only the presence of intact exendin, and negligible appearance of "daughter" peaks indicative of the buildup of degradation products. This is in contrast to other peptides studied (for example amylin and GLP-1) where the disappearance of the "parent" HPLC peak was associated with the appearance of "daughter" HPLC peaks, subsequently identified as subpeptide degradants. The absence of plasma degradants of exendin indicates little or no proteolysis at any site, including the renal circulation. Any clearance by the kidney, then, is via non-proteolytic means, namely filtration or active excretion (as occurs with para-amino hippurate).

Initial measurements of exendin clearance in man (120-130 mL/min), monkeys (~9 mL/min) and rats (3.2-4.4 mL/min) matched reported glomerular filtration rates in those species. To test whether renal filtration could be the principal mode of exendin elimination, studies were performed in overnight fasted nephrectomized male rats

infused with exendin at a constant rate. Steady-state plasma levels of exendin-4 were greatly increased in nephrectomized rats compared to rats with their kidneys intact. This data indicated that the kidney was responsible for at least 80% of the clearance of exendin 4 (see Figures 5 and 6). Exendin clearance rates in intact rats were, again, similar to glomerular filtration rates expected in those rats (4.2 mL/min). Taken together these results indicate that very little metabolism occurs systemically and that most of the clearance of exendin 4 is through the kidney via filtration (but not by renal intravascular proteolysis). The low amounts of immunoreactive full-length exendin in the urine are consistent with it being cleaved by brush border enzymes in the proximal tubule after filtration.

EXAMPLE 4 - EXENDIN-4 DECREASES GLUCAGON SECRETION DURING  
HYPERGLYCEMIC CLAMPS IN DIABETIC FATTY ZUCKER RATS

Absolute or relative hyperglucagonemia is often a feature of, for example, type 1 and type 2 diabetes mellitus, and the suppression of excessive glucagon secretion in these and other conditions described or referred to herein is a potential benefit of therapy using glucagonostatic agents. In this Example, the effect of exendin-4 on glucagon secretion in male anaesthetized Diabetic Fatty Zucker (ZDF) rats was examined. Using an hyperinsulinemic hyperglycemic clamp protocol, factors tending to influence glucagon secretion were held constant. Plasma glucose was clamped at ~34mM 60 min before beginning intravenous infusions of saline (n=7) or exendin-4 (0.21 $\mu$ g + 2.1 $\mu$ g/mL/h; n=7). Plasma glucagon concentration measured prior to these infusions were similar in both groups (306  $\pm$  30pM versus 252  $\pm$  32pM, respectively; n.s.).

Mean plasma glucagon concentration in exendin-4 infused rats was nearly half of that in saline-infused rats in the final 60 minutes of the clamp ( $165 \pm 18\text{pM}$  versus  $298 \pm 26\text{pM}$ , respectively;  $P < 0.002$ ). The hyperglycemic clamp protocol also enabled measurement of insulin sensitivity. Glucose infusion rate during the clamp was increased by  $111 \pm 7\%$  in exendin-4-treated versus control rats ( $P < 0.001$ ). In other words, exendin-4 exhibited a glucagonostatic effect in ZDF rats during hyperglycemic clamp studies, an effect that will be of therapeutic benefit in diabetic humans.

EXAMPLE 5 - METABOLIC EFFECTS OF EXENDIN-4 ON GLUCAGON  
SECRETION IN PEOPLE WITH TYPE 2 DIABETES

In this Example, the safety, tolerability, and efficacy of synthetic exendin-4 was evaluated in 8 male non-insulin using patients with type 2 diabetes who had discontinued other antidiabetic therapy for a minimum of 7 days. Each patient received subcutaneous (SC) injections of placebo (PBO) and 0.1, 0.2, and 0.3  $\mu\text{g/kg}$  exendin-4 48 hours apart in a single-blind, dose-rising, placebo controlled crossover design. Five patients also received a 0.4  $\mu\text{g/kg}$  dose. Plasma glucose, insulin and glucagon concentrations were assessed fasting and in response to a 7 Kcal/kg Sustacal® challenge administered at the time of exendin-4/PBO injection. Gastric emptying was evaluated by measuring serum acetaminophen concentrations following a 20 mg/kg oral dose of liquid acetaminophen administered with the Sustacal®. No safety issues were identified based upon reported adverse events, EKG and safety lab monitoring. Doses of 0.3 and 0.4  $\mu\text{g/kg}$  elicited a dose-dependent increase in nausea; vomiting occurred at the highest dose.

Plasma glucose concentrations were reduced in all doses of exendin-4 compared to PBO although insulin concentrations



were not significantly different. The 8 hour mean  $\pm$  SE changes in plasma glucose AUC from baseline were  $+391 \pm 187$ ,  $-263 \pm 108$ ,  $-247 \pm 64$ ,  $-336 \pm 139$ , and  $-328 \pm 70$  mg\*hr/dL for the PBO, 0.1, 0.2, 0.3, and 0.4  $\mu$ g/kg doses respectively. The 3 hr changes in plasma glucagon were  $+128.0 \pm 19.2$ ,  $-5.6 \pm 10.5$ ,  $-29.4 \pm 18.6$ ,  $-40.5 \pm 24.5$ , and  $+6.9 \pm 38.6$  pg\*hr/mL respectively. The gastric emptying rate was slowed in all doses and the mean total absorbed acetaminophen over 6 hours was reduced by 51%, 50%, 57% and 79% compared to PBO for 0.1, 0.2, 0.3, and 0.4  $\mu$ g/kg doses respectively. In summary, SC injection of exendin-4 to patients identified no safety issues, was tolerated at doses  $\leq 0.3$   $\mu$ g/kg, reduced plasma glucose and glucagon and slowed the rate of gastric emptying. These observations support the use of exendin for the treatment of conditions that would benefit from reduced glucagon levels and/or suppression of glucagon, including but not limited to type 1 and type 2 diabetes.

#### EXAMPLE 6: PEG MODIFIED EXENDIN 4

In the case of exendin 4, a 39 amino acid peptide with a molecular weight of 4187, modifications that increase its size and/or increase its anionic nature will decrease its ability to be filtered by the kidney. Because clearance of exendin 4 is largely by the kidney this will effectively increase its half life. Other properties of PEGylation (increased plasma half-life due to evasion of such renal and/or cellular clearance mechanisms that may exist; reduced immunogenicity and antigenicity; increased solubility; resistance to proteolysis; reduced toxicity (avoid dose spike); improved thermal and mechanical stability; improved permeability of the mucus or epithelial layer; and selective control over a specific biological function) are also of potential benefit for exendin 4 and exendin agonists.

In particular, because we have observed multiple pharmacologies (likely representing multiple receptor subtypes), different spectra of biological activities of exendin 4 may be selected by putting a PEG group at appropriate positions. Loss or alteration of bioactivity has been reported for PEGylated proteins which may be due to the presence of the PEG chains themselves, the particular site occupied by the PEG chain, or the coupling conditions having an adverse effect on the protein.

Primary considerations for PEG modification in terms of filtration at the kidney of exendin and exendin agonists are size and charge. Unmodified, exendin 4 has a molecular weight of approximately 4.2 kD and is anionic in nature with an overall net charge of approximately -2 at physiological pH. One, two or three PEG constituents may be covalently linked to exendin 4 or an analog of exendin 4, for example, with one PEG constituent being preferred. The size of the PEG can vary from a molecular weight of 500 to 20,000, preferably between 5,000 and 12,000.

Many of the methods for covalent attachment of PEG take advantage of the epsilon-amino group on lysine. Exendin 4 has two lysines that can be modified by attachment of PEG. An alanine scan of AC3177 (Leu<sup>14</sup>, Phe<sup>25</sup><sub>1-28</sub> exendin-4), a shortened analog of exendin 4, revealed positions that are sensitive to substitution by alanine. The two lysines at positions 12 and 27 were moderately affected by this substitution suggesting that loss of the lysine specific R group side chain (methylene chain plus epsilon-amino group) is tolerated. With regard to the full-length peptide, exendin 4, the two lysine positions are appropriate for PEG attachment (see compounds 1 and 2). In addition, depending on the chemistry used to conjugate the PEG, the epsilon-

amino groups at these positions may be masked thereby increasing the anionic nature of the peptide.

(201) HEGTFTSDLSK(PEG)QMEEEAVRLFIEWLKNGGPSSGAPPPS-NH<sub>2</sub>

(202) HEGTFTSDLSKQMEEEAVRLFIEWLK(PEG)NGGPSSGAPPPS-NH<sub>2</sub>

5 Based on the results of the alanine scan, other likely positions that may be modified by insertion of a Lys-PEG or equivalent, for example, are:

(203) HK(PEG)EGTFTSDLSKQMEEEAVRLFIEWLKNGGPSSGAPPPS-NH<sub>2</sub>

(204) HGEK(PEG)FTSDLSKQMEEEAVRLFIEWLKNGGPSSGAPPPS-NH<sub>2</sub>

10 (205) HEGTFTK(PEG)DLSKQMEEEAVRLFIEWLKNGGPSSGAPPPS-NH<sub>2</sub>

(206) HEGTFTSDK(PEG)SKQMEEEAVRLFIEWLKNGGPSSGAPPPS-NH<sub>2</sub>

(207) HEGTFTSDLK(PEG)KQMEEEAVRLFIEWLKNGGPSSGAPPPS-NH<sub>2</sub>

(208) HEGTFTSDLSKK(PEG)MEEAVRLFIEWLKNGGPSSGAPPPS-NH<sub>2</sub>

(209)\* HEGTFTSDLSKQMEK(PEG)EAVRLFIEWLKNGGPSSGAPPPS-NH<sub>2</sub>

15 (210)\* HEGTFTSDLSKQMEK(PEG)AVRLFIEWLKNGGPSSGAPPPS-NH<sub>2</sub>

(211) HEGTFTSDLSKQMEEEAK(PEG)RLFIEWLKNGGPSSGAPPPS-NH<sub>2</sub>

(212) HEGTFTSDLSKQMEEEAVRK(PEG)FIEWLKNGGPSSGAPPPS-NH<sub>2</sub>

(213)\* HEGTFTSDLSKQMEEEAVRLFIEK(PEG)WLKNGGPSSGAPPPS-NH<sub>2</sub>

(214) HEGTFTSDLSKQMEEEAVRLFIEK(PEG)LKNGGPSSGAPPPS-NH<sub>2</sub>

20 (215) HEGTFTSDLSKQMEEEAVRLFIEWLKK(PEG)GGPSSGAPPPS-NH<sub>2</sub>

The three positions\* above normally containing a glutamic acid that were indicated for modification with K(PEG) can also be modified by conjugation to the glutamic side chain carboxyl group, E(PEG).

25 Another analog in which the Lys-PEG can be added is at the supposed GlyGly turn:

(216) HEGTFTSDLSKQMEEEAVRLFIEWLKNK(PEG)GPSSGAPPPS-NH<sub>2</sub>

(217) HEGTFTSDLSKQMEEEAVRLFIEWLKNK(PEG)PSSGAPPPS-NH<sub>2</sub>

30 Positions 29-39 of exendin-4 may not be critical for the glucose lowering activity as evidenced by AC3177 having

nearly equipotent activity to exendin 4, and any of them, alone or in combination, can be substituted for K(PEG) or an equivalent.

5 One skilled in the art would readily appreciate that the present invention is well adapted to carry out the objects and obtain the ends and advantages mentioned, as well as those inherent therein. The molecular complexes and the methods, procedures, treatments, molecules, specific  
10 compounds described herein are presently representative of preferred embodiments are exemplary and are not intended as limitations on the scope of the invention. Changes therein and other uses will occur to those skilled in the art which are encompassed within the spirit of the invention are  
15 defined by the scope of the claims.

It will be readily apparent to one skilled in the art that varying substitutions and modifications may be made to the invention disclosed herein without departing from the scope and spirit of the invention.

20 All patents and publications mentioned in the specification are indicative of the levels of those skilled in the art to which the invention pertains. All patents and publications are herein incorporated by reference to the same extent as if each individual publication was  
25 specifically and individually indicated to be incorporated by reference.

The invention illustratively described herein suitably may be practiced in the absence of any element or elements, limitation or limitations, which is not specifically  
30 disclosed herein. Thus, for example, in each instance herein any of the terms "comprising", "consisting essentially of" and "consisting of" may be replaced with either of the other two terms. The terms and expressions

which have been employed are used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, but  
5 it is recognized that various modifications are possible within the scope of the invention claimed. Thus, it should be understood that although the present invention has been specifically disclosed by preferred embodiments and optional features, modification and variation of the concepts herein  
10 disclosed may be resorted to by those skilled in the art, and that such modifications and variations are considered to be within the scope of this invention as defined by the appended claims.

In addition, where features or aspects of the invention  
15 are described in terms of Markush groups, those skilled in the art will recognize that the invention is also thereby described in terms of any individual member or subgroup of members of the Markush group. For example, if X is described as selected from the group consisting of bromine,  
20 chlorine, and iodine, claims for X being bromine and claims for X being bromine and chlorine are fully described.

The invention has been described broadly and generically herein. Each of the narrower species and subgeneric groupings falling within the generic disclosure  
25 also form part of the invention. This includes the generic description of the invention with a proviso or negative limitation removing any subject matter from the genus, regardless of whether or not the excised material is specifically recited herein.

30 Other embodiments are within the following claims.

CLAIMS

1. A method of lowering plasma glucagon in a subject, comprising administering to said subject a therapeutically effective glucagon lowering amount of a compound selected from the group consisting of an exendin, an an exendin agonist, a modified exendin and a modified exendin agonist.
2. The method of claim 1 wherein said subject is suffering from necrolytic migratory erythema.
3. The method of claim 1 wherein said subject has a glucagonoma.
4. The method of any of claims 1-3 wherein said exendin agonist is an exendin.
5. The method of claim 4 wherein said exendin is exendin-4.
6. The method of any of claims 1-3 or 4 wherein said subject is a human.
7. The method of any of claims 1-3 wherein said modified exendin or exendin agonist is linked to one or more polyethylene glycol polymers.
8. The method of claim 7, wherein said one or more polyethylene glycol polymers each have molecular weights between 500 and 20,000.

His Ser Asp Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu  
 1 5 10 15  
 Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Pro Ser  
 20 25 30  
 Ser Gly Ala Pro Pro Ser-NH<sub>2</sub>  
 35

**Fig. 1**

His Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu  
 5 10 15  
 Glu Ala Val Arg Leu Phe Ile Glu Trp Leu Lys Asn Gly Pro Ser  
 20 25 30  
 Ser Gly Ala Pro Pro Ser-NH<sub>2</sub>  
 35

**Fig. 2**

1 Xaa<sub>1</sub> Xaa<sub>2</sub> Xaa<sub>3</sub> Gly Thr Xaa<sub>4</sub> Xaa<sub>5</sub> Xaa<sub>6</sub> Xaa<sub>7</sub> Xaa<sub>8</sub> Ser Lys Gln Xaa<sub>9</sub> Glu Glu Glu Ala Val Arg Leu  
 5 10 15 20  
 Xaa<sub>10</sub> Xaa<sub>11</sub> Xaa<sub>12</sub> Xaa<sub>13</sub> Leu Lys Asn Gly Gly Xaa<sub>14</sub> Ser Ser Gly Ala Xaa<sub>15</sub> Xaa<sub>16</sub> Xaa<sub>17</sub> Xaa<sub>18</sub>-Z  
 25 30 35

Compound [SEQ.ID.NO]	Xaa <sub>1</sub>	Xaa <sub>2</sub>	Xaa <sub>3</sub>	Xaa <sub>4</sub>	Xaa <sub>5</sub>	Xaa <sub>6</sub>	Xaa <sub>7</sub>	Xaa <sub>8</sub>	Xaa <sub>9</sub>	Xaa <sub>10</sub>	Xaa <sub>11</sub>	Xaa <sub>12</sub>	Xaa <sub>13</sub>	Xaa <sub>14</sub>	Xaa <sub>15</sub>	Xaa <sub>16</sub>	Xaa <sub>17</sub>	Xaa <sub>18</sub>	Z
1 [10]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
2 [11]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Ser	NH <sub>2</sub>	
3 [12]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Ser	NH <sub>2</sub>	
4 [13]	Tyr	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Ser	NH <sub>2</sub>	
5 [14]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Ser	NH <sub>2</sub>	
6 [15]	His	Gly	Asp	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Tyr	NH <sub>2</sub>	
7 [16]	His	Gly	Glu	Glu	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Ser	NH <sub>2</sub>	
8 [17]	His	Gly	Glu	Phe	Ser	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Ser	NH <sub>2</sub>	
9 [18]	His	Gly	Glu	Phe	Ser	Thr	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Ser	NH <sub>2</sub>	
10 [19]	His	Gly	Glu	Phe	Thr	Thr	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Ser	NH <sub>2</sub>	
11 [20]	His	Gly	Glu	Phe	Thr	Ser	Glu	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Ser	NH <sub>2</sub>	
12 [21]	His	Gly	Glu	Phe	Thr	Ser	Asp	pGly	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Ser	NH <sub>2</sub>	
13 [22]	His	Gly	Glu	Phe	Thr	Ser	Asp	pGly	Leu	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Ser	NH <sub>2</sub>	
14 [23]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	pGly	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Ser	NH <sub>2</sub>	
15 [24]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	pGly	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Ser	NH <sub>2</sub>	
16 [25]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	naph	Ile	Glu	Trp	Pro	Pro	Pro	Ser	NH <sub>2</sub>	
17 [26]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Val	Glu	Trp	Pro	Pro	Pro	Ser	NH <sub>2</sub>	

Fig. 3A



Compound [SEQ.ID.NO.]	Xaa <sub>1</sub>	Xaa <sub>2</sub>	Xaa <sub>3</sub>	Xaa <sub>4</sub>	Xaa <sub>5</sub>	Xaa <sub>6</sub>	Xaa <sub>7</sub>	Xaa <sub>8</sub>	Xaa <sub>9</sub>	Xaa <sub>10</sub>	Xaa <sub>11</sub>	Xaa <sub>12</sub>	Xaa <sub>13</sub>	Xaa <sub>14</sub>	Xaa <sub>15</sub>	Xaa <sub>16</sub>	Xaa <sub>17</sub>	Xaa <sub>18</sub>	Z
18 [27]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Val	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
19 [28]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
20 [29]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
21 [30]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Asp	Trp	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
22 [31]	His	Ala	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Phe	Pro	Pro	Pro	Pro	Ser	NH <sub>2</sub>
23 [32]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	tPro	tPro	tPro	tPro	Ser	NH <sub>2</sub>
24 [33]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	tPro	tPro	tPro	Ser	NH <sub>2</sub>
25 [34]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	hPro	hPro	hPro	hPro	Ser	NH <sub>2</sub>
26 [35]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	hPro	hPro	hPro	Ser	NH <sub>2</sub>
27 [36]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Phe	tPro	tPro	tPro	tPro	Ser	NH <sub>2</sub>
28 [37]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Phe	hPro	hPro	hPro	hPro	Ser	NH <sub>2</sub>
29 [38]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	MeAla	MeAla	MeAla	MeAla	Ser	NH <sub>2</sub>
30 [39]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Met	Phe	Ile	Glu	Trp	Pro	MeAla	MeAla	MeAla	Ser	NH <sub>2</sub>
31 [40]	His	Gly	Glu	Phe	Thr	Ser	Asp	Leu	Leu	Phe	Ile	Glu	Phe	MeAla	MeAla	MeAla	MeAla	Ser	NH <sub>2</sub>

Fig. 3B

Amino Acid Position	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Compound 1	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 2	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 3	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 4	His	Ala	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 5	His	Gly	Glu	Gly	Ala	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 6	His	Gly	Glu	Gly	Thr	Ala	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 7	His	Gly	Glu	Gly	Thr	Phe	Thr	Ala	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 8	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Ala	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 9	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ala	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 10	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Ala	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 11	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Ala	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 12	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Ala	Glu	Glu	Glu	Ala	Val	Arg
Compound 13	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Ala	Glu	Glu	Ala	Val	Arg
Compound 14	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Ala	Glu	Ala	Val	Arg
Compound 15	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Ala	Val	Arg
Compound 16	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Ala	Val	Arg
Compound 17	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Ala	Arg
Compound 18	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Ala
Compound 19	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 20	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg

Fig. 4A1

Amino Acid Position	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39
Compound 1	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	NH <sub>2</sub>								
Compound 2	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 3	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 4	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 5	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 6	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 7	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 8	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 9	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 10	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 11	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 12	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 13	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 14	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 15	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 16	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 17	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 18	Ala	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 19	Leu	Phe	Ile	Ala	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 20	Leu	Phe	Ile	Glu	Ala	Leu	Lys	Asn	NH <sub>2</sub>										

Fig. 4A2

09/899331

Amino Acid Position	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Compound 21	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 22	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 23	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 24	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 25	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 26	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 27	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 28	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 29	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 30	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 31	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 32	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 33	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 34	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 35	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 36	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 37	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 38	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 39	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 40	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 41	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg

Fig. 4A3

6/26

09/899331

Amino Acid Position	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39
Compound 21	Leu	Phe	Ile	Glu	Phe	Ala	Lys	Asn	NH2										
Compound 22	Leu	Phe	Ile	Glu	Phe	Leu	Ala	Asn	NH2										
Compound 23	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Ala	NH2										
Compound 24	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	Pro	Pro	Pro	NH2
Compound 25	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	Pro	Pro	Pro	NH2
Compound 26	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	Pro	Pro	NH2	
Compound 27	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	Pro	Pro	NH2	
Compound 28	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	Pro	NH2		
Compound 29	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	Pro	NH2		
Compound 30	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	NH2			
Compound 31	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	NH2			
Compound 32	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	NH2				
Compound 33	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	NH2				
Compound 34	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	NH2					
Compound 35	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	NH2					
Compound 36	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	NH2					
Compound 37	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	NH2					
Compound 38	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	NH2							
Compound 39	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	NH2							
Compound 40	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	NH2							
Compound 41	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	NH2									

Fig. 4A4

Amino Acid Position	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Compound 42	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 43	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 44	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 45	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 46	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 47	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 48	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 49	Arg	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 50	His	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 51	His	Gly	Glu	Gly	Thr	naph	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 52	His	Gly	Glu	Gly	Thr	Phe	Ser	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 53	His	Gly	Glu	Gly	Thr	Phe	Ser	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 54	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Glu	Leu	Ser	Lys	Gln	Met	Ala	Glu	Glu	Ala	Val	Arg
Compound 55	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	pGly	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 56	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 57	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 58	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 59	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Ala	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 60	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Ala	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 61	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Ala	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg

Fig. 4B1

8/26

Amino Acid Position	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39
Compound 42	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	NH2									
Compound 43	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	tPro	Ser	Ser	Gly	Ala	tPro	tPro	tPro	NH2
Compound 44	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	tPro	tPro	tPro	NH2
Compound 45	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Nme	Ser	Ser	Gly	Ala	Pro	Pro	NH2	
Compound 46	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Nme	Ser	Ser	Gly	Ala	Nme	Nme	NH2	
Compound 47	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	hPro	Ser	Ser	Gly	Ala	hPro	hPro	NH2	
Compound 48	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	hPro	Ser	Ser	Gly	Ala	hPro	NH2		
Compound 49	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	NH2			
Compound 50	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	NH2								
Compound 51	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2										
Compound 52	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH2										
Compound 53	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH2										
Compound 54	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH2										
Compound 55	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH2										
Compound 56	Leu	naph	Ile	Glu	Phe	Leu	Lys	Asn	NH2										
Compound 57	Leu	Phe	tBug	Glu	Trp	Leu	Lys	Asn	NH2										
Compound 58	Leu	Phe	Ile	Asp	Phe	Leu	Lys	Asn	NH2										
Compound 59	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	NH2					
Compound 60	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	NH2									
Compound 61	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	hPro	Ser	Ser	Gly	Ala	hPro	hPro	NH2	

Fig. 4B2

Compound  
No.

- 62 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu  
Phe Ile Glu Trp Leu Lys-NH<sup>E</sup>octanoyl Asn-NH<sub>2</sub>
- 63 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu  
Phe Ile Glu Phe Leu Lys-NH<sup>E</sup>octanoyl Asn-NH<sub>2</sub>
- 64 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu  
Phe Ile Glu Trp Leu Lys-NH<sup>E</sup>octanoyl Asn Gly Gly-NH<sub>2</sub>
- 65 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val  
Arg Leu Phe Ile Glu Phe Leu Lys-NH<sup>E</sup>octanoyl Asn Gly Gly-NH<sub>2</sub>
- 66 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu  
Phe Ile Glu Trp Leu Asn Lys-NH<sup>E</sup>octanoyl-NH<sub>2</sub>

10/26

SUBSTITUTE SHEET (RULE 26)

Fig. 4C



- Compound  
No.
- 67 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Ala Val Arg Leu  
Phe Ile Glu Phe Leu Asn Lys-NH<sup>E</sup> octanoyl-NH<sub>2</sub>
- 68 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu  
Phe Ile Glu Trp Leu Asn Lys-NH<sup>E</sup> octanoyl Gly Gly-NH<sub>2</sub>
- 69 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Ala Val Arg Leu  
Phe Ile Glu Phe Leu Asn Lys-NH<sup>E</sup> octanoyl Gly Gly-NH<sub>2</sub>

**Fig. 4D**

11/26

SUBSTITUTE SHEET (RULE 26)

Amino Acid Position	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Compound 70	Ala	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 71	His	Gly	Ala	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 72	His	Gly	Glu	Ala	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 73	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Ala	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 74	Ala	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 75	His	Gly	Ala	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 76	His	Gly	Glu	Ala	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 77	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Ala	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 78	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Ala	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 79	Ala	Ala	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 80	Ala	Ala	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 81	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 82	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 83	Ala	Gly	Asp	Gly	Ala	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 84	Ala	Gly	Asp	Gly	Ala	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 85	Ala	Gly	Asp	Gly	Thr	Nala	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 86	Ala	Gly	Asp	Gly	Thr	Nala	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 87	Ala	Gly	Asp	Gly	Thr	Phe	Ser	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 88	Ala	Gly	Asp	Gly	Thr	Phe	Ser	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 89	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ala	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg

Fig. 4E1

12/26

SUBSTITUTE SHEET (RULE 26)

09/899331

Amino Acid Position	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39
Compound 70	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 71	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 72	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 73	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 74	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 75	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 76	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 77	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 78	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 79	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 80	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 81	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 82	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 83	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 84	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 85	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 86	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 87	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 88	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 89	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										

Fig. 4E2

Amino Acid Position	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Compound 21	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ala	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 22	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Ala	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 23	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Ala	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 24	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Glu	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 25	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Glu	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 26	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Ala	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 27	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Ala	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 28	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Pgly	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 29	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Pgly	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 30	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ala	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 31	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ala	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 32	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Ala	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 33	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Ala	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 34	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Ala	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 35	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Ala	Leu	Glu	Glu	Glu	Ala	Val	Arg

Fig. 4E3

Amino Acid Position	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39
Compound 21	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 22	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 23	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 24	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 25	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 26	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 27	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 28	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 29	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 30	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 31	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 32	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 33	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 34	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 35	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										

Fig. 4E4

Amino Acid Position	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Compound 105	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Ala	Glu	Glu	Glu	Ala	Val	Arg
Compound 106	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Ala	Glu	Glu	Glu	Ala	Val	Arg
Compound 107	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	pGly	Glu	Glu	Glu	Ala	Val	Arg
Compound 108	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	pGly	Glu	Glu	Glu	Ala	Val	Arg
Compound 109	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Ala	Glu	Glu	Ala	Val	Arg
Compound 110	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Ala	Glu	Glu	Ala	Val	Arg
Compound 111	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Ala	Glu	Ala	Val	Arg
Compound 112	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Ala	Glu	Ala	Val	Arg
Compound 113	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Ala	Ala	Val	Arg
Compound 114	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Ala	Ala	Val	Arg
Compound 115	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Ala	Arg
Compound 116	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Ala	Arg
Compound 117	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Ala
Compound 118	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Ala
Compound 119	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 120	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 121	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 122	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 123	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 124	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg

Fig. 4F1

Amino Acid Position	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39
Compound 105	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 106	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 107	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 108	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 109	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 110	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 111	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 112	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 113	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 114	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 115	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 116	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 117	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 118	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 119	Ala	Phe	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 120	Ala	Phe	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 121	Leu	Nala	Ile	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 122	Leu	Nala	Ile	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 123	Leu	Phe	Val	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 124	Leu	Phe	Val	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										

Fig. 4F2

09/899331

Amino Acid Position	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Compound 125	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 126	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 127	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 128	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 129	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 130	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 131	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 132	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 133	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 134	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 135	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 136	Ala	Gly	Asp	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 137	Ala	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 138	His	Gly	Ala	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 139	His	Gly	Glu	Ala	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 140	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Ala	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg

Fig. 4F3



Amino Acid Position	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39
Compound 125	Leu	Phe	tGly	Glu	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 126	Leu	Phe	tGly	Glu	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 127	Leu	Phe	Ile	Asp	Trp	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 128	Leu	Phe	Ile	Asp	Phe	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 129	Leu	Phe	Ile	Glu	Ala	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 130	Leu	Phe	Ile	Glu	Ala	Leu	Lys	Asn	NH <sub>2</sub>										
Compound 131	Leu	Phe	Ile	Glu	Trp	Ala	Lys	Asn	NH <sub>2</sub>										
Compound 132	Leu	Phe	Ile	Glu	Phe	Ala	Lys	Asn	NH <sub>2</sub>										
Compound 133	Leu	Phe	Ile	Glu	Trp	Leu	Ala	Asn	NH <sub>2</sub>										
Compound 134	Leu	Phe	Ile	Glu	Phe	Leu	Ala	Asn	NH <sub>2</sub>										
Compound 135	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Ala	NH <sub>2</sub>										
Compound 136	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Ala	NH <sub>2</sub>										
Compound 137	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	Pro	Pro	Pro	NH <sub>2</sub>
Compound 138	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	Pro	Pro	Pro	NH <sub>2</sub>
Compound 139	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	Pro	Pro	Pro	NH <sub>2</sub>
Compound 140	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	Pro	NH <sub>2</sub>		

Fig. 4F4

Amino Acid Position	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Compound 141	Ala	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Ala	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 142	Ala	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 143	His	Gly	Ala	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 144	His	Gly	Glu	Ala	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 145	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Ala	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 146	Ala	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 147	His	Gly	Ala	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 148	His	Gly	Glu	Ala	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 149	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Ala	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 150	Ala	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg
Compound 151	His	Gly	Ala	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 152	His	Gly	Glu	Ala	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 153	His	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Ala	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 154	Ala	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 155	His	Gly	Ala	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 156	His	Gly	Ala	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 157	Ala	Gly	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Met	Glu	Glu	Glu	Ala	Val	Arg
Compound 158	Ala	Gly	Ala	Gly	Thr	Phe	Thr	Ser	Asp	Leu	Ser	Lys	Gln	Leu	Glu	Glu	Glu	Ala	Val	Arg

Fig. 4G1

Amino Acid Position	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39
Compound 141	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	Pro	NH2		
Compound 142	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	NH2			
Compound 143	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	NH2			
Compound 144	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	NH2				
Compound 145	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	NH2					
Compound 146	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	NH2						
Compound 147	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	NH2						
Compound 148	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	NH2							
Compound 149	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	NH2								
Compound 150	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	NH2									
Compound 151	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	tPro	Ser	Ser	Gly	Ala	tPro	tPro	tPro	NH2
Compound 152	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	tPro	tPro	tPro	NH2
Compound 153	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Nme	Ser	Ser	Gly	Ala	Nme	Nme	NH2	
Compound 154	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	hPro	Ser	Ser	Gly	Ala	hPro	NH2		
Compound 155	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	NH2			
Compound 156	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	NH2								
Compound 157	Leu	Phe	Ile	Glu	Trp	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	Pro	Pro	Pro	Ser
Compound 158	Leu	Phe	Ile	Glu	Phe	Leu	Lys	Asn	Gly	Gly	Pro	Ser	Ser	Gly	Ala	Pro	Pro	Pro	Ser

Fig. 4G2

Compound  
No.

159 4-Imidazolylpropionyl-Gly Glu Gly Thr Ser Ala Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu  
Phe Ile Glu Trp Leu Lys-NH<sup>E</sup>octanoyl Asn-NH<sub>2</sub>

160 4-Imidazolylpropionyl-Gly Glu Gly Thr Ser Ala Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu  
Phe Ile Glu Phe Leu Lys-NH<sup>E</sup>octanoyl Asn-NH<sub>2</sub>

161 4-Imidazolylpropionyl-Gly Glu Gly Thr Ser Ala Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu  
Phe Ile Glu Trp Leu Lys-NH<sup>E</sup>octanoyl Asn Gly Gly-NH<sub>2</sub>

162 4-Imidazolylpropionyl-Gly Glu Gly Thr Ser Ala Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu  
Phe Ile Glu Phe Leu Lys-NH<sup>E</sup>octanoyl Asn Gly Gly-NH<sub>2</sub>

163 4-Imidazolylpropionyl-Gly Glu Gly Thr Ser Ala Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu  
Phe Ile Glu Trp Leu Asn Lys-NH<sup>E</sup>octanoyl-NH<sub>2</sub>

164 4-Imidazolylpropionyl-Gly Glu Gly Thr Ser Ala Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu  
Phe Ile Glu Phe Leu Asn Lys-NH<sup>E</sup>octanoyl-NH<sub>2</sub>

**Fig. 4H**

22/26

SUBSTITUTE SHEET (RULE 26)

Compound  
No.

165 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Met Glu Glu Ala Val Arg Leu  
Phe Ile Glu Trp Leu Asn Lys-NH<sup>E</sup>octanoyl Gly Gly-NH<sub>2</sub>

166 4-Imidazolylpropionyl-Gly Glu Gly Thr Phe Thr Ser Ala Leu Ser Lys Gln Leu Glu Glu Ala Val Arg Leu  
Phe Ile Glu Phe Leu Asn Lys-NH<sup>E</sup>octanoyl Gly Gly-NH<sub>2</sub>

167 Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Ala Val Arg Leu Phe Ile Glu Trp  
Leu Lys-NH<sup>E</sup>octanoyl Asn -NH<sub>2</sub>

168 Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe  
Leu Lys-NH<sup>E</sup>octanoyl Asn -NH<sub>2</sub>

169 Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Ala Val Arg Leu Phe Ile Glu Trp  
Leu Lys-NH<sup>E</sup>octanoyl Asn Gly Gly-NH<sub>2</sub>

170 Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe  
Leu Lys-NH<sup>E</sup>octanoyl Asn Gly Gly-NH<sub>2</sub>

**Fig. 4I**

## Compound

No.


171 Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp  
Leu AsnLys-NH<sup>E</sup>octanoyl-NH<sub>2</sub>


172 Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe  
Leu Asn Lys-NH<sup>E</sup>octanoyl-NH<sub>2</sub>

173 Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Met Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Trp  
Leu Asn Lys-NH<sup>E</sup>octanoyl Gly Gly-NH<sub>2</sub>


174 Ala Gly Glu Gly Thr Phe Thr Ser Asp Leu Ser Lys Gln Leu Glu Glu Glu Ala Val Arg Leu Phe Ile Glu Phe  
Leu Asn Lys-NH<sup>E</sup>octanoyl Gly Gly-NH<sub>2</sub>

Fig. 4J

**Effect of functional nephrectomy on Exendin-4 clearance****Fig. 5**

**Fig. 6**





#3

## DECLARATION Utility Application

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled **METHODS FOR GLUCAGON SUPPRESSION** the specification of which

(Check One)

- ☐ is attached hereto OR  
☒ was filed on July 13, 2001 as United States Application Serial No. 09/889,331; based on PCT International Application No. PCT/US00/00942 filed January 14, 2001 and was amended on \_\_\_\_\_ (if applicable).

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment(s) referred to above.

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, § 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, § 119(a)-(d) or § 365(b) of any foreign application(s) for patent or inventor's certificate, or § 365(a) of any PCT international application which designated at least one country other than the United States of America, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or of any PCT international application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application Number(s)	Country	Date of Filing	Priority Claimed	
			Yes	No

I hereby claim the benefit under Title 35, United States Code §119(e) of any United States provisional application(s) listed below.

Application Number(s)	Filing Date
60/175,365	January 10, 2000
60/132,017	April 30, 1999
60/116,380	January 14, 1999

I hereby claim the benefit under Title 35, United States Code, § 120 of any United States application(s), or § 365(c) of any PCT international application designating the United States of America, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT international application in the manner provided by the first paragraph of Title 35, United States Code, § 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations § 1.56 which became available between the filing date of the prior application and the national or PCT international filing date of this application.

U.S. Parent Application Number	PCT Parent Number	Parent Filing Date	Status-Patented, Pending or Abandoned

I further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Title 18, United States Code, § 1001 and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

1-00

201	FULL NAME OF INVENTOR	FIRST Name <u>Andrew</u>	MIDDLE Initial <u>A.</u>	LAST Name <u>Young</u>	
	RESIDENCE & CITIZENSHIP	City <u>Point Loma</u> <u>La Jolla</u>	State or Foreign Country <u>California</u> <u>CA</u>	Country of Citizenship <u>New Zealand</u> <u>United States of America</u>	
	POST OFFICE ADDRESS	<u>P.O. Box 60501</u> <u>5679 LA JOLLA HERMOSA</u>	City <u>LA JOLLA</u> <u>Point Loma</u>	State or Country <u>California</u>	Zip Code <u>92037</u>
INVENTOR'S SIGNATURE <u>Andrew Young</u> DATE _____					


2-00

202	FULL NAME OF INVENTOR	FIRST Name <u>Bronislava</u>	MIDDLE Initial -	LAST Name <u>Gedulin</u>	
	RESIDENCE & CITIZENSHIP	City <u>San Diego</u>	State or Foreign Country <u>California</u> <u>CA</u>	Country of Citizenship <u>U.S.A.</u>	
	POST OFFICE ADDRESS	<u>14174 Half Moon Bay Drive</u>	City <u>San Diego</u>	State or Country <u>California</u>	Zip Code <u>92014</u>
INVENTOR'S SIGNATURE <u>B Gedulin</u> DATE <u>11/12/01</u>					

Rec'd PCT/PTO 18 DEC 2001

Patent

030639.0031.UTL1



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re the Application of:

Andrew Young, et al.

Serial No.: 09/889,331

International Application Filing Date:  
January 14, 2000

For: METHODS FOR GLUCAGON  
SUPPRESSION

)  
) Group Art Unit: To be assigned

)  
) Examiner: To be assigned

GRANT OF POWER OF ATTORNEY

Commissioner for Patents  
Washington, D.C. 20231

Dear Sir:

As representative of the assignee of an entire interest in the above-identified application, by virtue of an executed assignment of all inventors' rights in the application, the undersigned hereby appoints as attorney for the assignee herein, to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: John M. Benassi, Reg. No. 27,483; Richard

CERTIFICATE OF MAILING  
(37 C.F.R. §1.8a)

I hereby certify that this paper (along with anything referred to as being attached or enclosed) is being deposited with the United States Postal Service on the date shown below with sufficient postage as First Class Mail in an envelope addressed to the Commissioner for Patents, Washington, D.C. 20231.

November 13, 2001  
Date of Deposit

BARBARA J. KIELT  
Name of Person Mailing Paper

Barbara J. Kielt  
Signature of Person Mailing Paper


L. Blaylock, Reg. No. 43,503; Bradford J. Duft, Reg. No. 32,219; Jeffrey W. Guise, Reg. No. 34,613; Kurt M. Kjelland, Reg. No. 41,901; Stephen S. Korniczky, Reg. No. 34,853; Edward O. Kreusser, Reg. No. 38,523; Alma Levy, Reg. No. 43,751; F.T. Alexandra Mahaney, Reg. No. 37,668; Peter R. Munson, Reg. No. 43,821; Douglas C. Murdock, Reg. No. 37,549; Vicki G. Norton, Reg. No. 40,745; Douglas E. Olson, Reg. No. 22,798; Richard H. Pagliery, Reg. No. 44,276; John Peterson, Reg. No. 45,894; Jonathan Pettit, Reg. No. 40,770; Andrew D. Skale, Reg. No. 44,338; Franklin D. Ubell, Reg. No. 27,009; Peter D. Weinstein, Reg. No. 43,251; Anatoly Weiser, Reg. No. 43,229; Jessica R. Wolff, Reg. No. 37,261 and Lisa M. McGeehan, Reg. No. 41,185 of Brobeck, Phleger & Harrison LLP, 12390 El Camino Real, San Diego, California 92130; telephone (858) 720-2500 and and Molly A. Holman, Reg. No. 40,022 and Joanna Moore, Reg. No. 44,950, of Amylin Pharmaceuticals, Inc., 9373 Towne Centre Drive, San Diego, CA 92121, telephone (858) 731-1500.

In accordance with 37 C.F.R. § 3.73(b)(i), a copy of the executed assignment referenced above is submitted herewith. If any additional information is required, please advise Applicants accordingly.

Respectfully submitted,

AMYLIN PHARMACEUTICALS, INC.

Dated: 10/18/01


By: 

Name: Mark G. Foletta

Title: Vice President of Finance & CEO  
as Representative for Amylin  
Pharmaceuticals, Inc.

**AMYLIN PHARMACEUTICALS, INC.**  
9373 Towne Centre Drive  
San Diego, California 92121

United States Patent & Trademark Office  
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not present  
for scanning. (Document title)

☐ Page(s) \_\_\_\_\_ of \_\_\_\_\_ were not present  
for scanning. (Document title)

☒ *Scanned copy is best available. Drawings are dark*